

DISTANCES TO GALACTIC HIGH-VELOCITY CLOUDS. I. COHEN STREAM, COMPLEX GCP, CLOUD g1¹

B. P. WAKKER,² D. G. YORK,³ R. WILHELM,⁴ J. C. BARENTINE,⁵ P. RICHTER,⁶
T. C. BEERS,⁷ Ž. IVEZIĆ,⁸ AND J. C. HOWK⁹
Received 2007 July 31; accepted 2007 September 10

ABSTRACT

The high- and intermediate-velocity interstellar clouds (HVCs/IVCs) are tracers of energetic processes in and around the Milky Way. Clouds with near-solar metallicity about 1 kpc above the disk trace the circulation of material between disk and halo (the Galactic fountain). The Magellanic Stream consists of gas tidally extracted from the SMC, tracing the dark matter potential of the Milky Way. Several other HVCs have low metallicity and appear to trace the continuing accretion of infalling intergalactic gas. These assertions are supported by the metallicities (0.1 to 1 solar) measured for about 10 clouds in the past decade. Direct measurements of distances to HVCs have remained elusive, however. In this paper we present four new distance brackets, using VLT observations of interstellar Ca II H and K absorption toward distant Galactic halo stars. We derive distance brackets of 5.0 to 11.7 kpc for the Cohen Stream (likely to be an infalling low-metallicity cloud), 9.8 to 15.1 kpc for Complex GCP (also known as the Smith Cloud or HVC 40–15+100 and with still unknown origin), 1.0 to 2.7 kpc for an IVC that appears associated with the return flow of the fountain in the Perseus arm, and 1.8 to 3.8 kpc for cloud g1, which appears to be in the outflow phase of the fountain. Our measurements further demonstrate that the Milky Way is accreting substantial amounts of gaseous material, which influences the Galaxy’s current and future dynamical and chemical evolution.

Subject headings: Galaxy: halo — ISM: clouds — stars: distances — stars: horizontal-branch

1. INTRODUCTION

The evolution of galaxies is strongly driven by the gas in the interstellar medium (ISM). High- and low-mass stars form from gas in the disk and put back energy into the gas. Some gas escapes from the disk, forming a hot atmosphere. As this gas cools and rains back, it forms a “Galactic fountain.” The hot atmosphere has been called the “Galactic corona,” and it extends up to 10 kpc above the Galactic disk. There is also infall of new material that provides fuel for galaxy growth. This material can originate in small satellite galaxies, as gas tidally pulled out of passing galaxies, and from more pristine intergalactic gas. The cool clouds that fall in would be embedded in (or condense out of) a much more extended (100–200 kpc radius) hot corona, different from the one formed by star formation processes. Indirect evidence for infalling gas is provided by two arguments (Chiappini et al. 2001). First, at the current rate of star formation, all of the ISM would be turned into gas within about 1 Gyr. Second, the narrowness of the dis-

tribution of metallicities of long-lived stars is best explained if the metallicity of the ISM remains more or less constant over a Hubble time, which happens if there is an inflow of low-metallicity material from outside. The predicted inflow rate decreases with time and at present is on the order of $1 M_{\odot} \text{ yr}^{-1}$.

Direct observational evidence for the processes mentioned above is provided by the high-velocity clouds (see reviews by Wakker & van Woerden 1997; Richter 2006). They are defined observationally as clouds with velocities that are incompatible with differential galactic rotation. In practice, this means $|v_{\text{LSR}}| > 50 \text{ km s}^{-1}$. Historically, such gas has been divided into the high-velocity clouds (HVCs), having $|v_{\text{LSR}}| > 90$ or $> 100 \text{ km s}^{-1}$, and the intermediate-velocity clouds (IVCs), with $|v_{\text{LSR}}| = 50\text{--}100 \text{ km s}^{-1}$. They have been mapped extensively in 21 cm emission (see Hulsbosch & Wakker 1988; Wakker & van Woerden 1991; Hartmann & Burton 1997; Morras et al. 2000; Kalberla et al. 2005). The different origins of different clouds can be determined by measuring their metallicities and distances. Solar metallicity clouds near the disk trace the Galactic fountain. Distant (> 5 kpc), low-metallicity (0.01–0.2 solar) clouds trace infall. Long streams with intermediate metallicity are likely to be tidal. Intrinsic metallicities (Z) have been determined for a few HVCs (see van Woerden & Wakker 2004 for a summary). The best values are $Z \sim 0.15$ times solar for Complex C (Wakker et al. 1999; Richter et al. 2001b; Tripp et al. 2003; Sembach et al. 2004), while clouds in the Magellanic Stream have SMC-like metallicity (~ 0.25 solar; Lu et al. 1998; Gibson et al. 2000). Van Woerden & Wakker (2004) list approximate values for a total of eight HVCs, ranging from 0.1 to 1 times solar. In contrast, the IVCs are usually found to have near-solar metallicity (Wakker 2001; Richter et al. 2001a).

The main unknown for the HVCs remains their distances. If we knew these, we could derive empirical (rather than theoretical) estimates for the rate of accretion of low-metallicity (0.01–0.1 solar) gas, the rate of circulation of gas between disk and halo, and the mass of tidal material around the Milky Way. Furthermore, HVCs could be used to calibrate the fraction of ionizing photons

¹ Based on observations carried out at the European Southern Observatory (ESO), La Silla, under program ID 077.B-0359 (PI: P. Richter) with the UVES at the ESO Very Large Telescope, Paranal, Chile.

² Department of Astronomy, University of Wisconsin, Madison, WI 53706; wakker@astro.wisc.edu.

³ Astronomy and Astrophysics Center, University of Chicago, Chicago, IL 60637; don@oddjob.uchicago.edu.

⁴ Department of Physics and Astronomy, Texas Tech University, Lubbock, TX 79409; ron.wilhelm@ttu.edu.

⁵ Department of Astronomy, University of Texas, Austin, TX 78712; and Apache Point Observatory, Sunspot, NM 88349; jcb@astro.as.utexas.edu.

⁶ Institut für Physik, Universität Potsdam, 14469 Potsdam, Germany; prichter@astro.physik.uni-potsdam.de.

⁷ Department of Physics and Astronomy, Center for the Study of Cosmic Evolution (CSCE) and Joint Institute for Nuclear Astrophysics (JINA), Michigan State University, East Lansing, MI 48824; beers@pa.msu.edu.

⁸ Department of Astronomy, University of Washington, Seattle, WA 98195; ivezic@astro.washington.edu.

⁹ Department of Physics, University of Notre Dame, Notre Dame, IN 46556; jhowk@nd.edu.

escaping the Milky Way disk, as many HVCs are seen in H α emission (see Bland-Hawthorn & Maloney 1999).

Although the first paper on determining distances to HVCs was published four decades ago (Prata & Wallerstein 1967), progress has been slow. Wakker (2001) summarized the state of knowledge a few years back. It is clear that distances to several IVCs are on the order of 1–2 kpc. However, for HVCs Wakker (2001) only lists lower limits of 6 kpc to Complex C ($v_{\text{LSR}} \sim -150 \text{ km s}^{-1}$), 5 kpc to Complex H ($v_{\text{LSR}} \sim -180 \text{ km s}^{-1}$), and 0.3 kpc to the anticenter clouds (clouds with $v_{\text{LSR}} \sim -250 \text{ km s}^{-1}$ and $v_{\text{LSR}} \sim -100 \text{ km s}^{-1}$), as well as two upper limits: 10 kpc for the high-latitude part of Complex A ($v_{\text{LSR}} \sim -150 \text{ km s}^{-1}$), based on the RR Lyrae star AD UMa (van Woerden et al. 1999), and 13 kpc for Complex WE ($v_{\text{LSR}} \sim +110 \text{ km s}^{-1}$), using the O9.7 II star HD 156359 (Sembach et al. 1991). Since then, Wakker et al. (2003) have reported a lower limit of 8 kpc for Complex A, while Thom et al. (2006) used the star HE 1048+0231 to set an upper limit of 8.8 kpc to cloud WW 35 ($v_{\text{LSR}} \sim +100 \text{ km s}^{-1}$).

In this paper we report on our observations of 25 targets in the direction to two HVCs and two IVCs. These yield distance brackets to all four clouds. We describe the method used to derive the distances of these clouds in § 2. In § 3 we summarize our observations, while the manner in which we make our measurements is discussed in § 4. In § 5 we present the results for each of the four clouds, describing the distance determination, notes on each of the stars, a summary of Ca II/H I ratios, a discussion of the implied location in the Milky Way, and a discussion of their ionized hydrogen content. In § 6 we present a short general discussion of our results, while in § 7 we summarize the conclusions.

2. DETERMINING HVC/IVC DISTANCES

2.1. Basic Method

The distance to an HVC/IVC can be determined by looking for interstellar absorption at the velocity of the cloud in spectra of stars with known distances. A detection sets an upper limit, while a *significant* nondetection sets a lower limit. A significant nondetection means that the ratio of the expected equivalent width to the observed upper limit is sufficiently large (e.g., >10 ; the exact value depends on the ion that is observed and on knowledge of the cloud's ionic abundance; see Wakker 2001). The best kind of stars for this purpose are blue horizontal branch (BHB) and RR Lyrae stars (in their hot phase). This is because they are fairly numerous, their distances can be determined relatively well, and they are relatively blue, resulting in few confusing stellar lines. This method is described in detail by Wakker & van Woerden (1997) and Wakker (2001).

There are three steps in the method: (1) locate candidate probe stars; (2) obtain intermediate-resolution spectra and photometry to derive their distances, velocities, and abundances; and (3) obtain high spectral resolution (15 km s^{-1} or better) and high signal-to-noise ratio (S/N) spectra for stars at a range of distances; these stars should be selected to have minimal blending between stellar and interstellar lines. In this paper, we only summarize steps 1 and 2, while we report on the most extensive set of step 3 observations presented in the literature so far. Our extensive program covering steps 1 and 2 will be described in subsequent papers.

2.2. Stellar Probes

Until recently, most of the candidate stellar probes were found from three catalogs: the variable star lists of Kukarkin et al. (1969, 1970), the PG survey (Green et al. 1986), and the HK survey (Beers et al. 1996). These list many BHB and RR Lyrae candidates with distances up to about 5 kpc, as well as a few more

distant stars (up to ~ 10 kpc). The recent data from the Sloan Digital Sky Survey (SDSS) and the 2 Micron All Sky Survey¹⁰ (2MASS) now allow us to make further progress, adding candidate stars at up to 100 kpc distance.

The SDSS project and the creation of the archive are discussed by York et al. (2000). The details of the data products are described by Stoughton et al. (2002) and Adelman-McCarthy et al. (2007 and references therein). A 2.5 m telescope (Gunn et al. 2006) is equipped with two dual-beam spectrographs ($R \sim 1800$), operating between 3900 and 9000 Å, and with a CCD camera (Gunn et al. 1998) with a five-filter photometric system (Fukugita et al. 1996), calibrated on imaging nights with a 20 inch (0.5 m) Photometric Telescope (Hogg et al. 2001; Smith et al. 2002; Ivezić et al. 2004; Tucker et al. 2006). The CCD camera is used to scan the northern sky in the five filters, from which our RR Lyrae or BHB candidate stars are chosen, using color criteria (Sirko et al. 2004; Ivezić et al. 2005). The star positions are determined in the manner described by Pier et al. (2003). These stars have distances ranging from ~ 5 kpc up to 100 kpc, as estimated from their V magnitudes ($V = 0.56r + 0.44g + 0.056i$; Ivezić et al. 2000) and assuming an absolute magnitude of $M_V = 0.7$ for RR Lyrae stars (Layden et al. 1996) and $M_V = 1.040 + (B - V)\{-4.423 + (B - V)[17.74 + (B - V)(-35.73)]\}$ for BHB stars (Preston et al. 1991).

2MASS scanned the sky at $2''$ pixels, using automated 1.3 m telescopes on Mount Hopkins and the Cerro Tololo Inter-American Observatory, taking images at J , H , and K . It includes thousands of BHB candidates with distances up to ~ 8 kpc, which can be selected from their $J - H$ and $J - K$ colors (Brown et al. 2004).

For some stars the SDSS provides intermediate-resolution (~ 2 Å) spectra. To obtain similar spectra for more SDSS stars, as well as for the 2MASS-selected stars, we have a large observing project underway at the 3.5 m telescope at Apache Point Observatory (APO). Between 2003 and 2006 we obtained spectra at APO for some 200 stars.

Using these spectra, we can derive the effective temperature and gravity for each star, following the method outlined by Wilhelm et al. (1999). Combining with photometry, these parameters can be used to derive good distance estimates. We will describe these observations (as well as improvements to the original method) in a separate paper (R. Wilhelm et al. 2008, in preparation). Here we just present an outline.

Our method compares the observed photometry (either SDSS ugr or Johnson UBV) and observed spectroscopy. We compare the Balmer line widths and equivalent width of the Ca II K line with a grid of synthetic colors and spectroscopy parameters that are generated from Atlas9 model atmospheres. Combining the Balmer line widths with $u - g$ (or $U - B$) color indices allows the determination of $\log g$ values across the temperature range $6000 \text{ K} < T_{\text{eff}} < 12,000 \text{ K}$. The determined stellar parameters (T_{eff} , $\log g$, and $[\text{Fe}/\text{H}]$) are then used to estimate absolute magnitudes M_V by placing the star on the theoretical isochrones of Girardi et al. (2004). This method allows the determination of distances for stars of various metallicities on the zero-age horizontal branch, on the main sequence, as well as for terminal HB stars and post-main-sequence blue stragglers. We estimate errors in the stellar parameters by changing the input values for observables (i.e., line widths and colors) by $\pm 1 \sigma$ one observable at a time. We thus recalculate T_{eff} , $\log g$, $[\text{Fe}/\text{H}]$, M_V , and D many times. The error follows as the rms spread in the set of results for each parameter.

¹⁰ Vizie Online Data Catalog, 2246 (R. M. Cutri et al. 2003).

2.3. Implementation

The results of the programs described above yield a sample of stars with known distances in the direction of HVCs. These then need to be observed with high spectral resolution ($R > 20,000$), in order to separate the different interstellar absorption components, as well as to maximize the sensitivity to the $\sim 20 \text{ km s}^{-1}$ wide HVC/IVC absorptions. In principle, Mg II $\lambda\lambda 2796.352, 2803.531$ and Ca II $\lambda\lambda 1036.337, 1334.532$ are the best lines, as they saturate at low H I column densities ($\sim 7 \times 10^{17} \text{ cm}^{-2}$). However, they require sparsely available time on telescopes in space, which are not sensitive enough to observe many 16th to 18th magnitude BHB stars.

In the optical the strongest absorption lines are the Ca II $\lambda\lambda 3934.770, 3969.591$ K and H lines. Previous observations of Ca II absorption in HVCs with bright 21 cm emission have yielded six detections, giving Ca II/H I ratios of $(17 \pm 6) \times 10^{-9}$ for Complex A (Schwarz et al. 1995), $(22 \pm 1) \times 10^{-9}$ for Complex C (Wakker et al. 1996), $(3.3 \pm 0.9) \times 10^{-9}$ for Complex M (Bowen et al. 2000), $(33 \pm 4) \times 10^{-9}$ for the Magellanic Stream (Songaila & York 1980), $(66 \pm 8) \times 10^{-9}$ for the Leading Arm of the Magellanic Stream (West et al. 1985), and $(270 \pm 120) \times 10^{-9}$ for Complex WB (Robertson et al. 1991). There are also several Ca II detections toward extragalactic supernovae in directions with very faint or no 21 cm (d’Odorico et al. 1985, 1989; Meyer & Roth 1991; Vladilo et al. 1994; Ho & Filippenko 1995, 1996), which yield very high Ca II/H I ratios. Toward IVCs Ca II/H I ratios range from 8×10^{-9} to 55×10^{-9} in the IV Arch, from 12×10^{-9} to 17×10^{-9} in the LLIV Arch, 90×10^{-9} in Complex K, 50×10^{-9} in the PP Arch, and 65×10^{-9} in Complex gp (Wakker 2001).

Using these detections of high- and intermediate-velocity absorption (as well as HVC/IVC absorption in other ions, such as Fe II), Wakker & Mathis (2000) discovered that for Ca II, Fe II, etc. the $[\text{ion}]/\text{H I}$ ratio correlates with $N(\text{H I})$. The physical origin of this effect remains unexplained, as discussed by Wakker & Mathis (2000). For Ca II this correlation can be expressed as

$$R(\text{Ca II}) = \frac{N(\text{Ca II})}{N(\text{H I})} = -0.78 \{ \log[N(\text{H I}) - 19.5] \} - 7.76.$$

In fact, this relation is a better predictor of the Ca II/H I ratio than the assumption that $R(\text{Ca II})$ is constant in a particular cloud. Therefore, we use this relation to convert the known H I column densities in HVCs and IVCs in the direction of a candidate target star to predict the Ca II equivalent width. This allows us to construct a sample of stars for each HVC by selecting the ones with the largest predicted equivalent widths in a set of distance intervals (e.g., 1–3 kpc, 3–6 kpc, 6–9 kpc, etc.). We have done this for 60 HVC/IVC fields. In this paper we report on observation of stars in three fields, covering two HVCs and two IVCs.

Previous authors have set distance brackets or limits to HVCs/IVCs using the nominal distances to stars that give upper limits and significant lower limits. However, the errors in the stellar distances have been ignored in the past. In this paper we take them into account. Since the error distribution is centered on the nominal distance, the probability that the upper limit is less than the *nominal* distance to the star is 50%. If the error distribution of the stellar distances were Gaussian, the probability that the upper limit is less than $D_{\text{nom}} + 0.47\sigma(D)$ is 68%. Thus, the 68% (i.e., 1σ) confidence interval on the upper distance limit to the HVC is given by $D_{\text{nom}} + 0.47\sigma(D)$. A similar reasoning provides the lower limit of the bracket. At present, we do not calculate the actual error distri-

bution for the distance, so for convenience we assume that it is Gaussian. For future work we will improve this step.

3. OBSERVATIONS

3.1. Stellar Data

For the 2006 April–September semester we were given an allocation of 37 hr on ESO’s Very Large Telescope (VLT) to observe 24 stars and one QSO (proposal ID 077.B-0359). For our observations we used the UVES installed on the VLT together with a $1''$ slit and the blue arm (CD2). This setup provides a wavelength coverage from ~ 3730 to 4990 \AA , a pixel size of 0.03 \AA ($\sim 2.3 \text{ km s}^{-1}$), and a spectral resolution of $R \sim 40,000$, corresponding to an FWHM of $\sim 7.5 \text{ km s}^{-1}$. During our observations the seeing typically was $\leq 0.8''$. The raw data were reduced with the standard UVES pipeline, which includes flat-fielding, bias and sky subtraction, and a relative wavelength calibration. To compare with the 21 cm data, the final spectra were then shifted to an LSR velocity scale. For multiple integrations of the same background source, the individual spectra were co-added, weighted by their inverse variances. In the continuum the S/N per resolution element in the UVES data typically ranges from about 40 to 100.

The three HVCs that we selected for our VLT observations have in common that they are above the horizon during the southern winter. They are the “Cohen Stream,” “Complex GCP,” and “cloud g1.” Most stars in the Cohen Stream field also lie projected onto another, more nearby IVC. We describe these clouds and the stars projected on them in the next section. Figure 1 shows the locations of these three clouds on a map of the HVC sky. The map also shows the three stars toward which HVC absorption was found previously.

Table 1 presents the stellar data. Coordinates are defined in the names for 2MASS and SDSS stars and given in the notes for the remaining stars. The two “BS” stars are from the HK survey (Beers et al. 1996), and the numbers represent plate number and star ID on that plate. Seventeen stars in the VLT sample have classification spectra from our APO observations, and one spectrum is from the SDSS (see col. [2] of the table). We also obtained photometry for most stars, with the source of the data listed in column (3) and the resulting magnitudes in columns (4), (5), and (6). The extinction (from Schlegel et al. 1998) is given in column (7). For five stars we can use SDSS photometry. For 12 stars (11 from 2MASS, 1 from the HK survey) we obtained this photometry ourselves. For seven stars we do not have photometry data. Three of these are 2MASS stars, so we used the star’s B magnitude from the 2MASS catalog. For the three named RR Lyrae stars we used the photometry listed in the Kukarkin et al. (1970) catalog. Finally, Strömberg photometry was obtained by Moehler et al. (1990) for PG 0142+148.

For all stars we give the stellar parameters derived directly from the high-resolution spectra, while for the 17 stars with intermediate-resolution spectroscopy we give additional lines showing the stellar parameters resulting from our APO data, for comparison. The stellar parameters are derived using the photometry, the Balmer lines, Ca II K and H, and Ca I $\lambda 4227.918$. A full description of this method will be given by R. Wilhelm et al. (2008, in preparation). Columns (8) to (11) list the resulting effective temperature, gravity, metallicity, and stellar type, respectively. Combining the stellar parameters with photometry and isochrones, we can derive the star’s distance, given in column (13).

The VLT data were used to derive the stellar velocities listed in column (12). This was done by fitting a Gaussian to the stellar Ca II, Ca I, and Fe I lines, then averaging the fitted central velocities. For some of the stars the stellar velocity varies between




FIG. 1.—All-sky map of the HVC sky, based on the data of Hulsbosch & Wakker (1988) and Morras et al. (2000). Colors represent LSR velocities, as coded in the wedge. Contour levels are at brightness temperatures of 0.05, 0.25, 0.5, and 1 K. The positions of the clouds discussed in this paper are shown by the thick circles. Note that cloud g1 does not show up in this map because its velocity ($+70 \text{ km s}^{-1}$) falls outside the velocity range covered by Hulsbosch & Wakker (1988). Also shown are the locations of the three stars toward which high-velocity absorption was previously detected.

exposures (see the notes to the table). In these cases, we did not use all of the exposures, but only the ones with the least confusing separation between stellar and interstellar lines. The listed velocity is valid for that spectrum.

There are often differences in the temperature, gravity, and metallicity derived from the VLT spectrum when compared to those values derived from the APO spectra. When the differences are more than 1σ , they are noted in the following subsection.

3.2. Distance Estimation

3.2.1. Nonproblematic Stars

For eight stars (2MASS J014936.48+143914.6, 2MASS J021409.31+090105.3, 2MASS J195927.29+000822.3, 2MASS J195823.36–002719.0, 2MASS J195912.00–002645.3, 2MASS J213327.04+133026.7, 2MASS J213520.04+133045.0, and BS 17578–0016), the determination of the stellar distance based on the VLT and APO spectra is the same to within 15%. For four stars (2MASS J014825.88+132305.3, SDSS J014843.61+130411.5, 2MASS J195922.75+000300.0, and 2MASS J213451.58+134017.5) the determinations differ by 20%–40%. Since they are still the same within $\sim 1 \sigma$, the results are still compatible, just relatively inaccurate.

In the case of 2MASS J213451.58+134017.5 the metallicity indicated by the Ca II K line is about 1 dex less than that found from spectral regions with many metallic lines (mostly Fe). The star is probably a main-sequence star with solar abundance, so we adopt a nominal $\log g = 4.0$ to derive the distance. The lower resolution of the APO data makes that result less reliable, so we are not worried by the $\sim 1 \sigma$ difference in the nominal distance.

For one star (2MASS J195925.49–000519.1) the difference between the distance derived from the high- and the intermediate-resolution spectrum is larger than the nominal error ($1.5 \pm 0.2 \text{ kpc}$ vs. $2.6 \pm 0.4 \text{ kpc}$). However, these values are still somewhat consistent, and this is not an important star for our program to determine HVC distances. We therefore did not try to correct this discrepancy.

For two stars (SDSS J015735.65+135254.2 and BS 17578–0015) we have no APO classification spectrum, but we do have

photometry, and we are confident that the resulting classifications and distances are good.

Two stars are too hot (i.e., $T > 12,000 \text{ K}$) for our method, but distances are given in the literature. These are PG 0142+148, which was analyzed by Moehler et al. (1990), and PG 2134+125, which is the central star of the planetary nebula NGC 7094. Phillips (2004) summarizes distance determinations for planetary nebulae and includes NGC 7094.

3.2.2. Problematic Stars (Non-RR Lyrae)

For two stars the distances derived from the VLT and APO spectra differ substantially.

The VLT spectrum of 2MASS J021651.34+080150.2 shows many stellar lines, which blend in the APO spectrum, making it difficult to determine the gravity. In addition, we do not have photometry for this star, so that we cannot directly fit T_{eff} and $\log g$ (see R. Wilhelm et al. 2008, in preparation). We instead use a grid of spectral models with different temperatures and gravities and find the best match. As a result of these problems, the APO spectrum suggests low gravity, while the VLT spectrum indicates high gravity. This leads to a dichotomy in the classification; we have more confidence in the lower (VLT based) distance.

The VLT spectrum of 2MASS J195741.61–004009.7 also shows many stellar lines. In fact, the line density is such that it explains why the APO spectrum of this star is noisy, making its analysis difficult. The determination of the star's gravity is so uncertain that the APO data suggest that it is a BHB star, implying $D = 11.1 \pm 3.3 \text{ kpc}$, while the VLT spectrum indicates a main-sequence F star at $0.7 \pm 0.6 \text{ kpc}$. We judge that it is more likely to be nearby than to be distant.

3.2.3. RR Lyrae Stars

The final five stars are RR Lyrae stars, a classification based on the derived stellar parameters, but also on the fact that we can see the stellar lines shift between different VLT exposures. To estimate a distance for an RR Lyrae star requires a determination of its average magnitude. We usually do not have phase information, so we do not know whether the VLT spectrum or the SDSS photometry

TABLE 1
STELLAR DATA

Object (1)	Tel. (2)	Phot. (3)	$y/V/r$ (4)	B/g (5)	U/u (6)	E_{B-V} (7)	T_{eff} (K) (8)	$\log g$ (cm s^{-2}) (9)	$\log Z$ (Z_{\odot}^{-1}) (10)	Type (11)	$v(*)$ (km s^{-1}) (12)	D (kpc) (13)
PG 0142+148 ^a	VLT	Str	13.73	0.048	26200	5.10	...	sdB	-74 ± 2	1.2 ± 0.5
	APO	Str	13.73	0.048	26200	5.10	...	sdB	-67	1.2 ± 0.5
2MASS J014825.88+132305.3.....	VLT	UBV	13.89	14.06	14.01	0.071	7927 ± 140	4.08 ± 0.78	-1.99 ± 0.12	AV	-167 ± 2	2.1 ± 1.3
	APO	UBV	13.89	14.06	14.01	0.071	8120 ± 141	4.47 ± 0.63	-1.70 ± 0.12	AV	-155	1.2 ± 0.7
2MASS J021651.34+080150.2.....	VLT	2MS	16.80	0.102	6250 ± 250	4.50 ± 0.50	-1.00 ± 0.25	FV	213 ± 1	2.2 ± 1.0
	APO	2MS	16.80	0.102	6250 ± 250	2.75 ± 0.50	-1.25 ± 0.25	FHB	209	21.0 ± 8.6
2MASS J021409.31+090105.3.....	VLT	UBV	13.55	13.79	13.95	0.092	7744 ± 46	3.47 ± 0.17	-1.39 ± 0.16	AV	-37 ± 1	3.3 ± 0.8
	APO	UBV	13.55	13.79	13.95	0.092	7646 ± 76	3.25 ± 0.12	-1.71 ± 0.09	FHB	-2	3.1 ± 0.2
	APO	UBV	13.55	13.79	13.95	0.092	7749 ± 115	3.53 ± 0.16	-1.14 ± 0.14	AV	-25	3.3 ± 0.6
2MASS J014936.48+143914.6.....	VLT	UBV	14.05	14.21	14.27	0.044	8113 ± 106	3.42 ± 0.27	-0.03 ± 0.06	AV	-66 ± 7	4.6 ± 0.8
	APO	UBV	14.05	14.21	14.27	0.044	7901 ± 233	3.34 ± 0.37	-0.81 ± 0.28	FHB/AV	-30	4.0 ± 1.0
SDSS J014843.61+130411.5.....	VLT	SDSS	14.20	14.59	15.84	0.084	5880 ± 112	3.00 ± 0.25	-0.66 ± 0.11	FV	-57 ± 2	5.0 ± 1.2
	APO	UBVS	14.43	15.02	15.07	0.084	5774 ± 100	2.75 ± 1.07	-0.89 ± 0.32	FHB	-97	6.3 ± 2.5
SDSS J020033.50+141154.0.....	VLT	SDSS	15.48	15.58	16.89	0.059	6644 ± 22	2.00 ± 0.00	-1.32 ± 0.18	RR	-29 ± 3 ^b	8.7 ± 1.3
	SDSS	UBVS	15.58	15.87	15.97	0.059	6911 ± 21	2.15 ± 0.09	-1.73 ± 0.21	RR	8	8.7 ± 1.3
SDSS J015133.91+141105.2.....	VLT	SDSS	16.64	16.89	17.94	0.049	6823 ± 312	3.87 ± 0.37	-1.90 ± 0.22	RR	-190 ± 2 ^b	11.2 ± 1.0
	APO	UBVS	16.81	17.25	17.15	0.049	6530 ± 24	3.12 ± 0.09	-1.73 ± 0.30	AV	-176	11.2 ± 1.0
SDSS J015735.65+135254.2.....	VLT	SDSS	17.92	17.80	19.01	0.061	8109 ± 156	3.10 ± 0.13	-2.98 ± 0.15	FHB	1 ± 2	30.4 ± 3.9
	VLT	UBVS	17.92	17.99	18.01	0.061	8109 ± 156	3.10 ± 0.13	-2.98 ± 0.15	FHB	1 ± 2	30.4 ± 3.9
SDSS J014631.99+133506.3.....	VLT	SDSS	17.23	17.14	17.35	0.057	QSO	0.69 ^c	...
	VLT	UBVS	17.26	17.32	16.60	0.057	QSO	0.69 ^c	...
2MASS J195741.61-004009.7.....	VLT	2MS	16.56	16.90	...	0.240	6250 ± 250	4.50 ± 0.50	-1.00 ± 0.25	FV	73 ± 2	0.7 ± 0.6
	APO	2MS	16.56	16.90	...	0.240	6250 ± 250	3.25 ± 0.25	-1.25 ± 0.25	FHB	^d	11.1 ± 3.3
2MASS J195927.29+000822.3.....	VLT	UBV	14.88	15.28	15.39	0.214	7755 ± 115	4.52 ± 0.25	-1.01 ± 0.09	AV	-1 ± 3	1.1 ± 0.5
	APO	UBV	14.88	15.28	15.39	0.214	7463 ± 257	4.41 ± 0.35	-0.84 ± 0.26	AV	-35	1.1 ± 0.5
2MASS J195925.49-000519.1.....	VLT	UBV	14.24	14.70	14.93	0.215	7660 ± 45	4.18 ± 0.09	-0.14 ± 0.07	AV	34 ± 2	1.5 ± 0.2
	APO	UBV	14.24	14.70	14.93	0.215	7233 ± 124	3.67 ± 0.22	-0.09 ± 0.08	AV	-61	2.6 ± 0.4
2MASS J195912.00-002645.3.....	VLT	UBV	13.96	14.42	14.52	0.156	7124 ± 92	3.73 ± 0.04	-0.95 ± 0.06	AV	24 ± 4	1.8 ± 0.3
	APO	UBV	13.96	14.42	14.52	0.156	7089 ± 88	4.03 ± 0.22	-0.02 ± 0.06	AV	27	1.7 ± 0.4
2MASS J195922.75+000300.0.....	VLT	UBV	17.40	18.18	18.33	0.213	5762 ± 81	3.00 ± 0.50	0.00 ± 0.11	FHB	21 ± 2	9.0 ± 3.7
	APO	UBV	17.40	18.18	18.33	0.213	5750 ± 250	3.00 ± 0.50	-0.75 ± 0.25	FHB/AV	16	11.9 ± 3.9
V1172 Aql ^e	VLT	Kuk	16.5	0.182	6500 ± 250	2.50 ± 0.50	-0.85 ± 0.25	RR	-122 ± 1 ^b	10.5 ± 1.4
2MASS J195823.36-002719.0.....	VLT	UBV	16.85	17.57	17.78	0.203	5911 ± 91	2.50 ± 0.50	-0.35 ± 0.13	FHB	40 ± 2	13.1 ± 3.2
	APO	UBV	16.85	17.57	17.78	0.203	6164 ± 42	2.39 ± 0.31	-1.01 ± 0.13	FHB	26	11.0 ± 3.0
V1084 Aql ^f	VLT	Kuk	17.0	0.148	7000 ± 250	2.50 ± 0.50	-1.22 ± 0.25	RR	-32 ± 2	14.5 ± 1.3
BS 17578-0015 ^g	VLT	GSC	10.19	10.45	...	0.110	7125 ± 250	2.50 ± 0.25	-0.50 ± 0.25	FHB	-9 ± 5	1.2 ± 0.3
2MASS J213520.04+133045.0.....	VLT	UBV	13.22	13.44	13.74	0.131	7733 ± 58	4.00 ± 0.17	-0.72 ± 0.15	Am	26 ± 6	1.3 ± 0.2
	APO	UBV	13.22	13.44	13.74	0.131	7794 ± 180	4.00 ± 0.30	-1.55 ± 0.08	Am	-5 ^b	1.5 ± 0.5
2MASS J213451.58+134017.5.....	VLT	UBV	13.62	13.90	14.16	0.106	7473 ± 171	4.00 ± 0.25	-0.00 ± 0.02	Am	22 ± 4	2.0 ± 0.6
	APO	UBV	13.62	13.90	14.16	0.106	7365 ± 17	4.00 ± 0.17	-0.72 ± 0.15	Am	-10	1.3 ± 0.3
PG 2134+125 ^h	VLT	UBV	17.02	17.45	17.08	0.131	>40000	CSPN	-55 ± 9	2.2 ± 0.8
BS 17578-0016 ⁱ	VLT	UBV	12.59	12.60	12.42	0.114	11772 ± 280	3.83 ± 0.10	0.00	AV	-35 ± 5	3.6 ± 0.5
	APO	UBV	12.59	12.60	12.42	0.114	12000 ± 200	4.00 ± 0.25	0.00 ± 0.25	AV	-53	3.1 ± 0.8
FW Peg ^j	VLT	Kuk	14.0	0.120	6500 ± 250	2.50 ± 0.50	-0.82 ± 0.25	RR	-225 ± 2 ^b	3.8 ± 0.8
2MASS J213327.04+133026.7.....	VLT	UBV	14.98	15.02	14.90	0.111	11285 ± 176	3.97 ± 0.08	0.00	AV	-117 ± 3	8.2 ± 0.9
	APO	UBV	14.98	15.02	14.90	0.111	11500 ± 200	4.25 ± 0.25	0.00 ± 0.25	AV	-118	7.2 ± 0.4

NOTES.—Col. (1): Object name. Col. (2): Source of spectroscopic data: APO=Apache Point Observatory; VLT=Very Large Telescope. Col. (3): Source of photometry data: SDSS=Sloan Digital Sky Survey *ugr*; UBVS=SDSS photometry converted to *UBV*; *UBV*=our own *UBV* photometry; Str=Strömgren photometry; GSC=value from *HST* Guide Star Catalogue; Kuk=value from Kukarkin et al. (1970). Cols. (4)–(6): Magnitude: *U*, *B*, *V* when col. (3) has “UBV;” SDSS *r*, *g*, *u* when col. (3) has “SDSS;” Strömgren *y* when col. (3) has “Str;” *V* magnitude from Kukarkin et al. (1970) when col. (3) has “Kuk.” Col. (7): E_{B-V} . Col. (8): Derived effective temperature. Col. (9): Derived gravity. Col. (10): Derived abundance, relative to solar. Col. (11): Spectral type: FHB=field horizontal branch; RR=RR Lyrae; AV, FV=main-sequence A or F star. Col. (12): Derived stellar velocity; formal errors were not derived for velocities measured in the APO spectra: the pixels are about 50 km s⁻¹ wide. Col. (13): Derived distance; we prefer the distance derived using the VLT spectrum; for RR Lyrae stars we estimate the average magnitude, as explained in the text.

^a Temperature, gravity, and distance from Mochler et al. (1990); star located at (R.A., decl.) = (01^h45^m39.5^s, +15°04′42″).

^b Stellar velocity is variable; the listed velocity is valid for the spectrum in which the interstellar lines are searched for.

^c Redshift instead of velocity given for this QSO.

^d Too many stellar lines to reliably determine stellar velocity in low-resolution spectrum.

^e Star located at (R.A., decl.) = (20^h02^m23.0^s, -00°32′09″).

^f Star located at (R.A., decl.) = (20^h03^m43.5^s, +00°53′38″).

^g Star located at (R.A., decl.) = (21^h34^m17.8^s, +13°38′04″).

^h Distance from Phillips (2004); this is the central star of the planetary nebula NGC 7094; star located at (R.A., decl.) = (21^h36^m52.2^s, +13°35′00″).

ⁱ Star located at (R.A., decl.) = (21^h34^m45.4^s, +13°52′50″).

^j Star located at (R.A., decl.) = (21^h32^m59.0^s, +12°52′24″).

was taken during maximum, minimum, or somewhere in between. Moreover, the magnitudes given by Kukarkin et al. (1970) for V1084 Aql, V1172 Aql, and FW Peg are photographic, not *UBV*. They are also quantized, in steps of ~ 0.25 mag. Therefore, we use the following reasoning to estimate the distances to the five RR Lyrae stars in our sample.

We start by looking at SDSS J015133.91+141105.2. The SDSS photometry of this star gives $V = 16.81$ (using the transformation from *ugriz* to *UBV* given by Ivezić et al. 2000). We have six VLT exposures of this star, taken 28 and 16 days apart, with one run of four exposures in four consecutive hours. The star’s flux varies between 6.2×10^{-16} and 20.5×10^{-16} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$, giving 13.3×10^{-16} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$ on average, corresponding to $V = 16.14$. Since the range in flux is about 1.3 mag, it is reasonable to conclude that 16.1 ± 0.2 is a good estimate for the average magnitude of this star. Combined with an absolute magnitude for RR Lyrae stars of $M_V = 0.7$ (Layden et al. 1996) and extinction $A_V = 0.16$ (Schlegel et al. 1998), this implies a distance of 11.2 ± 1.0 kpc.

The flux of SDSS J020033.49+141153.9 varies from 10.7×10^{-16} to 21.6×10^{-16} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$ between the three spectra of this star. The average flux is about 16×10^{-16} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$, or $V \sim 16.0$. The temperature that is derived is about 6600 K, which suggests that the star is in its cool phase. The SDSS equivalent V magnitude is 15.58. These results are compatible if the SDSS observation caught it near average while the VLT observation caught it near minimum. Based on this, we set the average magnitude to $V = 15.6 \pm 0.3$. With $A_V = 0.19$ this implies $D = 8.7 \pm 1.3$ kpc.

For V1084 Aql Kukarkin et al. (1970) gave $\langle P \rangle = 17.0$. Bond (1978) included this star in a sample of RR Lyrae stars for which he obtained photometry. He finds $\langle V \rangle = 17.0$, with an accuracy of about 0.2 mag. The star’s flux varies from 7.4×10^{-16} to 14.5×10^{-16} ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$ in our four VLT exposures. On average this flux corresponds to $V = 16.35$. We derive a temperature of 7000 K, which suggests that we observed the star near maximum phase, which is compatible with the relatively high flux and the relatively low number of stellar lines. Using $A_V = 0.49$ (Schlegel et al. 1998) then implies a distance of 14.5 ± 1.3 kpc.

Kukarkin et al. (1970) give an average magnitude for V1172 Aql of 16.5 ± 0.25 . We find an average flux in the VLT spectra that corresponds to $V = 16.36$. The temperature that is derived from the three VLT exposures is about 6500 K, suggesting that we observed it in a cooler, fainter phase. We thus assume that the Kukarkin et al. (1970) magnitude is slightly low and calculate the distance using $\langle V \rangle = 16.4 \pm 0.3$, which for $A_V = 0.60$ gives $D = 10.5 \pm 1.4$ kpc.

The star FW Peg has $\langle V \rangle = 14.0$ in Kukarkin et al. (1970). The VLT spectrum gives a flux corresponding to $V = 14.68$ and a temperature of 6500 K. This is compatible if we caught the star near minimum. We thus use $\langle V \rangle = 14.0 \pm 0.5$ to calculate its distance. With $A_V = 0.40$ this gives 3.8 ± 0.8 kpc.

4. MEASUREMENTS

We now summarize the measurements we made on the stars and then discuss for each cloud in turn our previous knowledge, followed by a more general discussion of our results concerning their distances, Ca II/H I ratios, and location in the Milky Way.

4.1. Stellar Measurements

Table 2 presents the stellar data that pertain to searching for the interstellar lines. We give the star’s Galactic location in columns (2) and (3) and repeat the distance from Table 1. Column (5) gives the total VLT exposure times (but see the notes to the table

for exposures that were excluded for the final analysis). We also determined the star’s continuum flux, as well as the flux and S/N at the velocity of the HVCs (cols. [6], [7], and [8]). This was done by fitting a polynomial to the line-free regions of the continuum. To calculate the flux at the HVC’s velocity, the stellar Ca II absorption was considered as “continuum,” since we are interested in the flux and S/N appropriate for measuring the interstellar line against the background of the stellar flux. The order of the fitted polynomial lies between 1 and 4. The S/N is calculated per 7 km s $^{-1}$ resolution element, i.e., calculated after rebinning the spectrum to 3 pixels. It is the ratio of the flux to the rms of the fit around the line-free regions of the spectrum used in the fit. This value is compared with the nominal error in the flux to make sure that it is similar.

4.2. H I and Ca II Measurements

Table 3 lists the Ca II interstellar measurements. The table contains the following columns. Columns (1) and (2) repeat the stellar name and distance. Columns (3) and (4) give the velocity and the logarithm of the H I column density of the HVC, IVC, or low-velocity gas in the sight line. The H I 21 cm emission data were taken from the Leiden-Argentina-Bonn Survey (LAB; Kalberla et al. 2005). This survey provides spectra on a $0.5^\circ \times 0.5^\circ$ grid, with a $36'$ beam. We calculated a weighted average of all spectra that lie within 0.6° from each star as the best approximation to the H I spectrum toward each star. To determine $v(\text{H I})$ and $N(\text{H I})$, we fitted Gaussians to these spectra, using one Gaussian component for the high- or intermediate-velocity emission and between two and four for the low-velocity gas. It should be noted that the intermediate- and high-velocity components usually are not exactly Gaussian, so the fit gives only an approximation to $N(\text{H I})$.

The uncertainty in $N(\text{H I})$ is dominated by the fact that the Dwingeloo telescope has a large beam on the sky. Wakker et al. (2001) showed that the value of $N(\text{H I})$ measured with a $9'$ beam can be up to a factor of 2.5 lower or higher than the value measured with a $35'$ beam. Further, when comparing measurements of $N(\text{H I})$ made with a $9'$ beam with those made with a $1'-2'$ beam, the latter are between 75% and 125% of the former. The origin of these differences is clearly small-scale structure in the cloud, including edge effects if the beam is near a cloud’s edge. The distribution of the ratio $N(\text{H I, real})/N(\text{H I, LAB})$ has an average of about 1 and an rms of about 0.2. As a result, we assume a 20% error for $N(\text{H I})$ when calculating Ca II/H I ratios. Ideally, we would like to obtain high-resolution ($1'$ or better) H I 21 cm data toward each of our targets, but we do not have such data at this time. Even then, there is always the possibility of a hole in the distribution of the H I in the direction of the star, and a nondetection would still not imply that the star lies in front of the HVC. However, we use the apparent hierarchical structure of the H I in HVCs (see Schwarz & Wakker 2004) to argue that it is possible to account for the uncertainty in $N(\text{H I})$ by using a “safety factor,” as described below under the discussion of column (13).

Columns (5) and (6) give the predicted K and H equivalent widths, respectively. These are derived from a predicted Ca II column density, combined with the observed line width of the H I emission. The prediction for the column density is derived from the correlation between $N(\text{Ca II})$ and $N(\text{H I})$ discovered by Wakker & Mathis (2000), but with a cloud-dependent correction factor. The correction factor is found by comparing the nominal prediction [given a value of $N(\text{H I})$] for the Ca II/H I ratio with the actual observed value and then averaging the resulting quotient between all detections found for each cloud. We discuss the correction factors in § 5, but note here that the quotient of nominal prediction and actual value tends to be a constant value in a particular cloud, although it varies between clouds.

TABLE 2
OBSERVATIONAL DATA

Object (1)	Longitude (deg) (2)	Latitude (deg) (3)	D (kpc) (4)	T_{exp} (ks) (5)	Flux (cont) (10^{-16} ergs cm^{-2} s^{-1} \AA^{-1}) (6)	S/N (cont) (7)	Flux (HVC) (10^{-16} ergs cm^{-2} s^{-1} \AA^{-1}) (8)	S/N (HVC) (9)
Cohen Stream								
PG 0142+148.....	141.87	-45.79	1.2 ± 0.5	0.5	226	84	226	78
2MASS J014825.88+132305.3.....	143.56	-47.19	2.1 ± 1.3	1.0	137	83	128	80
2MASS J021651.34+080150.2.....	156.43	-49.21	2.2 ± 1.0	26.1 ^a	4.2	85	3.5	79
2MASS J021409.31+090105.3.....	154.81	-48.67	3.3 ± 0.8	2.7 ^b	190	84	132	55
2MASS J014936.48+143914.6.....	143.36	-45.90	4.6 ± 0.8	1.3	114	80	54	38
SDSS J014843.61+130411.5.....	143.81	-47.47	5.0 ± 1.2	3.0	45	96	15	42
SDSS J020033.50+141154.0.....	147.14	-45.40	8.7 ± 1.3	8.0 ^c	22	43	11	28
SDSS J015133.91+141105.2.....	144.23	-46.19	11.2 ± 1.0	21.1 ^d	17	99	13	69
SDSS J015735.65+135254.2.....	146.36	-45.96	30.4 ± 3.9	15.0 ^e	4.0	37	4.0	36
SDSS J014631.99+133506.3.....	142.83	-47.15	***	14.0	7.9	51	7.9	51
Complex GCP								
2MASS J195741.61-004009.7.....	40.04	-15.01	0.7 ± 0.6	8.4 ^f	5.6	37	0.8	7.6
2MASS J195927.29+000822.3.....	41.00	-15.02	1.1 ± 0.5	1.6	38	52	17	35
2MASS J195925.49-000519.1.....	40.79	-15.12	1.5 ± 0.2	1.0	67	42	10	14
2MASS J195912.00-002645.3.....	40.43	-15.24	1.8 ± 0.3	1.0	84	55	22	25
2MASS J195922.75+000300.0.....	40.91	-15.05	9.0 ± 3.7	1.7	2.5	11	0.5	3.3
V1172 Aql.....	40.74	-15.98	10.5 ± 1.4	8.2 ^g	11	52	7.6	52
2MASS J195823.36-002719.0.....	40.32	-15.07	13.1 ± 3.2	5.6 ^h	3.9	24	0.5	4.6
V1084 Aql.....	42.21	-15.60	14.5 ± 1.3	11.0 ⁱ	10	55	8.6	47
Cloud g1								
BS 17578-0015.....	67.04	-27.17	1.2 ± 0.3	0.5	2.5	77	1.9	70
2MASS J213520.04+133045.0.....	67.12	-27.44	1.3 ± 0.2	1.5	160	81	59	34
2MASS J213451.58+134017.5.....	67.17	-27.25	2.0 ± 0.6	2.6 ^j	84	101	41	47
PG 2134+125.....	66.78	-28.20	2.2 ± 0.8	3.0	249	103	241	148
BS 17578-0016.....	67.33	-27.10	3.6 ± 0.5	0.5	658	106	603	101
FW Peg.....	66.15	-27.39	3.8 ± 0.8	3.0 ^k	67	71	64	73
2MASS J213327.04+133026.7.....	66.78	-27.10	8.2 ± 0.9	3.0 ^l	40	73	40	91

NOTES.— Col. (1): Object name. Cols. (2) and (3): Galactic longitude and latitude of the star. Col. (4): Distance estimate from Table 1; asterisks mean that the distance to this QSO is off the scale for the column. Col. (5): Total exposure time (but see footnotes). Col. (6): Stellar flux in the continuum. Col. (7): Signal-to-noise ratio in the continuum. Col. (8): Stellar flux at the velocity of the HVC. Col. (9): Signal-to-noise ratio at the velocity of the HVC.

^a There are nine exposures: since the stellar velocity is constant, they are all combined.

^b There are four exposures of this star, but for unknown reasons the flux in the second is much lower, so we exclude it.

^c There are three exposures; the stellar velocity is -2 km s^{-1} in the first exposure, -8 km s^{-1} in the second, -29 km s^{-1} in the third. We only use the third exposure, as in exposures 1 and 2 a stellar line is shifted on top of the interstellar Ca II K line.

^d This star appears to be binary: in four of the seven exposures the stellar lines are at -190 km s^{-1} (and we combine these four), but in the remaining three exposures the stellar line is much broader, appears to have two components, and is centered at -178 , -138 , and -139 km s^{-1} ; it also confuses the interstellar Ca II K.

^e There are five exposures, with constant stellar velocity; however, in the first exposure a spike (probably a cosmic ray) confuses the interstellar K line.

^f The first of the three exposures has lower stellar flux, so we do not use it.

^g The stellar velocity changed from -128 to -121 to -115 km s^{-1} over the three contiguous hours of observing; since there are no stellar lines interfering with the interstellar Ca II K, we averaged the three exposures.

^h The two exposures of this star were combined.

ⁱ The stellar velocity changed from -36 to -31 to -26 km s^{-1} over the four contiguous hours of observing; since there are no stellar lines interfering with the interstellar Ca II K, we averaged these.

^j The two exposures of this star were combined.

^k The stellar velocity is -225 km s^{-1} in the first exposure, -170 km s^{-1} in the second; the wide stellar Ca II K line extends across the interstellar line in the second exposure, halving the flux, so we only use the first exposure.

^l The two exposures of this star were combined.

Columns (7) and (8) list the measured K and H line equivalent width detections or 3σ limits. For detections the absorption lines were integrated over the range of velocities where the flux lies below the continuum, which is $20\text{--}25 \text{ km s}^{-1}$ wide for the Cohen Stream, the IVC projected onto the Cohen Stream, and Complex GCP and about 40 km s^{-1} wide for cloud g1. The equivalent width limits for nondetections were determined by integrating the spectrum over a 20 km s^{-1} velocity range and setting the upper limit to 3 times the 1σ error in the equivalent width. The equivalent

width error includes the uncertainty associated with the noise, as well as the uncertainty associated with continuum placement.

In columns (9) and (10) we give the Ca II column density derived by integrating the apparent optical depth profile (Savage & Sembach 1991):

$$N = \int N_a(v) dv = \frac{m_e c}{\pi e^2} f \lambda \ln \frac{C(v)}{F(v)},$$

where $C(v)$ is the continuum and $F(v)$ is the observed flux.

TABLE 3
INTERSTELLAR ABSORPTION LINE RESULTS

Object (1)	D (kpc) (2)	v_{LSR} (km s^{-1}) (3)	$\log N(\text{H I})$ (4)	W_{exp} (K) ($\text{m}\text{\AA}$) (5)	W_{exp} (H) ($\text{m}\text{\AA}$) (6)	W_{obs} (K) ($\text{m}\text{\AA}$) (7)	W_{obs} (H) ($\text{m}\text{\AA}$) (8)	$\log N$ (K) (9)	$\log N$ (H) (10)	A (K) (10^{-9}) (11)	A (H) (10^{-9}) (12)	U, L (13)
Cohen Stream ($D = 5.0\text{--}11.7$ kpc)												
PG 0142+148.....	1.2 ± 0.5	-105	19.26	20	10	<7	<8	<10.88	<11.38	<4.2	<13.2	L
2MASS J014825.88+132305.3.....	2.1 ± 1.3	-106	19.41	22	11	<6	<15	<10.92	<11.38	<3.2	<9.3	L
2MASS J021651.34+080150.2.....	2.2 ± 1.0	-112	19.54	24	12	<10	<9	<11.05	<11.37	<3.2	<6.8	L
2MASS J021409.31+090105.3.....	3.3 ± 0.8	-110	19.08	19	9
2MASS J014936.48+143914.6.....	4.6 ± 0.8	-106	19.30	21	10	<9	<11	<11.02	<11.48	<5.2	<15	L
SDSS J014843.61+130411.5.....	5.0 ± 1.2	-103	19.18	19	10	<10	<10	<11.08	<11.48	<7.9	<20	...
SDSS J020033.50+141154.0.....	8.7 ± 1.3	-108	18.94	17	9	<17	<19	<11.28	<11.72	<22	<60	...
SDSS J015133.91+141105.2.....	11.2 ± 1.0	-104	19.26	20	10	17 ± 3	<28	11.33 ± 0.07	<11.54	11.7 ± 2.9	<19	U
SDSS J015735.65+135254.2.....	30.4 ± 3.9	-104	18.90	17	8	10 ± 3	<18	11.06 ± 0.20	<11.63	14.5 ± 7.6	<54	U
SDSS J014631.99+133506.3.....	***	-105	19.51	23	11	17 ± 3	11 ± 3	11.31 ± 0.09	11.41 ± 0.12	6.3 ± 1.7	7.9 ± 2.5	U
IV-South ($D = 1.0\text{--}2.7$ kpc)												
PG 0142+148.....	1.2 ± 0.5	-51	19.54	28	14	<7	<8	<10.94	<11.25	<2.5	<5.1	L
2MASS J014825.88+132305.3.....	2.1 ± 1.3	-41	19.11	23	11	25 ± 2	...	11.50 ± 0.04	...	24 ± 5	...	U
2MASS J021651.34+080150.2.....	2.2 ± 1.0	-75	18.93	21	10
2MASS J021409.31+090105.3.....	3.3 ± 0.8	-63	18.77	19	9
2MASS J014936.48+143914.6.....	4.6 ± 0.8	-51	19.49	27	14	26 ± 4	<14	11.50 ± 0.06	<11.50	10.2 ± 2.5	<10.2	U
SDSS J014843.61+130411.5.....	5.0 ± 1.2	-35	19.18	23	12	<12	<10	<11.02	<11.49	<6.9	<20	...
SDSS J020033.50+141154.0.....	8.7 ± 1.3	-46	18.86	20	10
SDSS J015133.91+141105.2.....	11.2 ± 1.0	-49	19.23	24	12	16 ± 3	26 ± 2	11.29 ± 0.08	11.50 ± 0.04	11.5 ± 3.0	19 ± 4	U
SDSS J015735.65+135254.2.....	30.4 ± 3.9	-48	18.95	21	10	22 ± 3	<21	11.42 ± 0.10	<11.80	30 ± 9	<71	U
SDSS J014631.99+133506.3.....	***	-43	19.20	24	12	30 ± 3	19 ± 3	11.55 ± 0.06	11.66 ± 0.10	22 ± 5	29 ± 8	U
Complex GCP ($D = 9.8\text{--}15.1$ kpc)												
2MASS J195741.61-004009.7.....	0.7 ± 0.6	89	20.16	61	32	<43	<40	<11.70	<12.19	<3.5	<10.7	...
2MASS J195927.29+000822.3.....	1.1 ± 0.5	93	20.19	64	33	<13	<18	<11.17	<11.61	<1.0	<2.6	L
2MASS J195925.49-000519.1.....	1.5 ± 0.2	93	20.15	62	32	<28	...	<11.58	...	<2.7
2MASS J195912.00-002645.3.....	1.8 ± 0.3	93	20.13	62	32	<15	<16	<11.18	<11.49	<1.1	<2.3	L
2MASS J195922.75+000300.0.....	9.0 ± 3.7	93	20.17	63	32	<124	<158	<12.36	<12.54	<16	<23	...
V1172 Aql.....	10.5 ± 1.4	94	20.03	59	30	<11	<13	<11.13	<11.56	<1.3	<3.4	L
2MASS J195823.36-002719.0.....	13.1 ± 3.2	90	20.12	61	31	<68	<57	<12.09	<12.25	<9.3	<13.5	...
V1084 Aql.....	14.5 ± 1.3	91	20.08	61	31	51 ± 3	37 ± 10	11.82 ± 0.02	11.96 ± 0.13	5.5 ± 1.1	7.6 ± 2.4	U
Cloud g1 ($D = 1.8\text{--}3.8$ kpc)												
BS 17578-0015.....	1.2 ± 0.3	71	19.04	74	39	<5	<8	<10.73	<11.02	<4.9	<9.5	L
2MASS J213520.04+133045.0.....	1.3 ± 0.2	71	19.04	75	39	<8	<11	<10.99	<11.59	<8.9	<36	L
2MASS J213451.58+134017.5.....	2.0 ± 0.6	71	18.95	72	38	<8	<10	<10.94	<11.35	<9.8	<25	L
PG 2134+125.....	2.2 ± 0.8	70	18.92	69	37	<4	<5	<10.66	<11.00	<5.5	<12.0	L
BS 17578-0016.....	3.6 ± 0.5	67	18.51	55	29	30 ± 2	14 ± 3	11.56 ± 0.03	11.52 ± 0.14	112 ± 24	102 ± 35	U
	3.6 ± 0.5	80	18.08	42	23	44 ± 2	31 ± 3	11.74 ± 0.02	11.87 ± 0.04	457 ± 94	617 ± 137	U
FW Peg.....	3.8 ± 0.8	67	19.23	82	43	127 ± 4	73 ± 4	12.25 ± 0.01	12.26 ± 0.03	105 ± 21	107 ± 23	U
2MASS J213327.04+133026.7.....	8.2 ± 0.9	72	19.11	77	41	66 ± 3	27 ± 9	11.92 ± 0.02	11.88 ± 0.14	65 ± 13	59 ± 20	U

TABLE 3—Continued

Object (1)	D (kpc) (2)	v_{LSR} (km s^{-1}) (3)	$\log N(\text{H I})$ (4)	W_{exp} (K) ($\text{m}\text{\AA}$) (5)	W_{exp} (H) ($\text{m}\text{\AA}$) (6)	W_{obs} (K) ($\text{m}\text{\AA}$) (7)	W_{obs} (H) ($\text{m}\text{\AA}$) (8)	$\log N$ (K) (9)	$\log N$ (H) (10)	A (K) (10^{-9}) (11)	A (H) (10^{-9}) (12)	U, L (13)
Low-Velocity Gas near CS												
PG 0142+148.....	1.2 ± 0.5	−3	20.56	203	112	94 ± 2	65 ± 4	12.12 ± 0.01	12.22 ± 0.03	3.6 ± 0.7	4.6 ± 1.0	U
2MASS J014825.88+132305.3.....	2.1 ± 1.3	−6	20.65	208	115	98 ± 3	64 ± 7	12.16 ± 0.01	12.23 ± 0.05	3.2 ± 0.6	3.8 ± 0.9	U
2MASS J021651.34+080150.2.....	2.2 ± 1.0	−12	20.81	226	127	115 ± 6	<11	12.24 ± 0.02	<12.23	2.7 ± 0.6	<2.6	U
2MASS J021409.31+090105.3.....	3.3 ± 0.8	−13	20.75	213	119
2MASS J014936.48+143914.6.....	4.6 ± 0.8	−4	20.59	206	114	124 ± 5	52 ± 7	12.26 ± 0.02	12.13 ± 0.07	4.7 ± 1.0	3.5 ± 0.9	U
SDSS J014843.61+130411.5.....	5.0 ± 1.2	−5	20.69	216	121	120 ± 5	78 ± 4	12.26 ± 0.02	12.31 ± 0.04	3.7 ± 0.8	4.2 ± 0.9	U
SDSS J020033.50+141154.0.....	8.7 ± 1.3	−4	20.68	212	118	120 ± 18	102 ± 16	12.30 ± 0.08	12.49 ± 0.06	4.2 ± 1.1	6.5 ± 1.6	U
SDSS J015133.91+141105.2.....	11.2 ± 1.0	−6	20.62	204	113	183 ± 3	123 ± 7	12.57 ± 0.01	12.61 ± 0.03	8.9 ± 1.8	9.8 ± 2.0	U
SDSS J015735.65+135254.2.....	30.4 ± 3.9	−6	20.66	213	118
SDSS J014631.99+133506.3.....	***	−4	20.47	193	106	155 ± 4	76 ± 4	12.35 ± 0.02	12.30 ± 0.03	7.6 ± 1.5	6.8 ± 1.5	U
Low-Velocity Gas near GCP												
2MASS J195741.61−004009.7.....	0.7 ± 0.6	32	20.47	177	101	181 ± 16	<74	12.50 ± 0.06	<12.35	10.7 ± 2.5	<7.6	U
2MASS J195927.29+000822.3.....	1.1 ± 0.5	30	20.43	176	101	186 ± 6	89 ± 9	12.52 ± 0.02	12.41 ± 0.05	12.3 ± 2.5	9.5 ± 2.1	U
2MASS J195925.49−000519.1.....	1.5 ± 0.2	32	20.40	176	101
2MASS J195912.00−002645.3.....	1.8 ± 0.3	35	20.37	170	97	124 ± 10	84 ± 9	12.24 ± 0.05	12.35 ± 0.05	7.4 ± 1.7	9.5 ± 2.2	U
2MASS J195922.75+000300.0.....	9.0 ± 3.7	32	20.42	174	99	366 ± 46	<222	12.91 ± 0.15	<12.84	31 ± 16	<26	U
V1172 Aql.....	10.5 ± 1.4	27	20.41	175	100	230 ± 4	152 ± 6	12.58 ± 0.01	12.67 ± 0.02	14.8 ± 3.0	18 ± 4	U
2MASS J195823.36−002719.0.....	13.1 ± 3.2	32	20.40	175	100	303 ± 24	250 ± 22	12.87 ± 0.13	12.96 ± 0.07	30 ± 12	36 ± 9	U
V1084 Aql.....	14.5 ± 1.3	32	20.33	167	95	240 ± 3	130 ± 11	12.65 ± 0.01	12.57 ± 0.04	21 ± 4	17 ± 4	U
Low-Velocity Gas near GCP												
2MASS J195741.61−004009.7.....	0.7 ± 0.6	0	20.82	200	117	118 ± 11	94 ± 13	12.31 ± 0.06	12.46 ± 0.06	3.1 ± 0.7	4.4 ± 1.1	U
2MASS J195927.29+000822.3.....	1.1 ± 0.5	0	20.85	202	118	213 ± 5	121 ± 7	12.61 ± 0.02	12.56 ± 0.03	5.8 ± 1.2	5.1 ± 1.1	U
2MASS J195925.49−000519.1.....	1.5 ± 0.2	0	20.82	202	118
2MASS J195912.00−002645.3.....	1.8 ± 0.3	0	20.79	199	116	154 ± 9	111 ± 8	12.42 ± 0.01	12.50 ± 0.03	4.3 ± 0.9	5.1 ± 1.1	U
2MASS J195922.75+000300.0.....	9.0 ± 3.7	0	20.84	207	122	193 ± 43	<160	12.60 ± 0.31	<12.71	5.8 ± 3.2	<7.4	U
V1172 Aql.....	10.5 ± 1.4	1	20.78	196	114	128 ± 5	182 ± 5	12.32 ± 0.02	12.78 ± 0.02	3.5 ± 0.7	10.0 ± 2.0	U
2MASS J195823.36−002719.0.....	13.1 ± 3.2	0	20.80	203	119	200 ± 18	150 ± 17	12.67 ± 0.16	12.73 ± 0.07	7.4 ± 3.5	8.5 ± 2.2	U
V1084 Aql.....	14.5 ± 1.3	1	20.82	199	117	248 ± 4	181 ± 9	12.67 ± 0.01	12.74 ± 0.02	7.1 ± 1.4	8.3 ± 1.7	U
Low-Velocity Gas near g1												
BS 17578−0015.....	1.2 ± 0.3	2	20.82	221	123	115 ± 3	74 ± 3	12.25 ± 0.01	12.30 ± 0.02	2.7 ± 0.5	3.0 ± 0.6	U
2MASS J213520.04+133045.0.....	1.3 ± 0.2	4	20.84	228	128	235 ± 3	176 ± 3	12.66 ± 0.01	12.74 ± 0.01	6.6 ± 1.3	7.9 ± 1.6	U
2MASS J213451.58+134017.5.....	2.0 ± 0.6	4	20.82	220	123	155 ± 9	108 ± 6	12.39 ± 0.02	12.48 ± 0.02	3.7 ± 0.8	4.6 ± 0.9	U
PG 2134+125.....	2.2 ± 0.8	1	20.84	230	129	178 ± 2	107 ± 2	12.46 ± 0.00	12.47 ± 0.01	4.2 ± 0.8	4.3 ± 0.9	U
BS 17578−0016.....	3.6 ± 0.5	2	20.79	219	123	216 ± 2	132 ± 3	12.58 ± 0.00	12.58 ± 0.01	6.2 ± 1.2	6.2 ± 1.3	U
FW Peg.....	3.8 ± 0.8	3	20.79	197	116	287 ± 3	197 ± 5	12.72 ± 0.01	12.78 ± 0.01	8.5 ± 1.7	9.8 ± 2.0	U
2MASS J213327.04+133026.7.....	8.2 ± 0.9	2	20.80	226	127	260 ± 2	...	12.65 ± 0.00	...	7.1 ± 1.4	...	U

NOTES.—Col. (1): Object name. Col. (2): Distance; asterisks mean that the distance to this QSO is off the scale for the column. Col. (3): Velocity of HVC or IVC in the direction of the star. Col. (4): Logarithm of 21 cm H I column density in the HVC. Cols. (5)–(8): Expected and observed equivalent width of the HVC Ca II absorption. Cols. (9) and (10): Logarithm of integrated apparent Ca II column density (in cm^{-2}) based on the K or H line profile. Cols. (11) and (12): Ca II/H I ratio, using $N(\text{K})$ or $N(\text{H})$; the error includes a 20% error in the H I column density. Col. (13): Conclusion drawn concerning HVC or IVC distance from this sight line: U means upper limit, L means lower limit, and no entry is given if the nondetection is not significant.

TABLE 4
SUMMARY OF CLOUD PARAMETERS

Quantity	Cohen Stream	Perseus Arm IVC	Complex GCP	Cloud g1
Distance (kpc)	5.0–11.7	1.0–2.7	9.8–15.1	1.8–3.8
Radius (kpc)	11.5–15.7	9.1–10.2	6.2–9.7	8.0–7.8
Height (kpc).....	3.7–8.6	0.7–1.8	2.5–3.9	0.8–1.7
Size (deg).....	1 × 12	3 × 8	2 × 8	2 × 2
H I Parameters				
$N(\text{H I})$ (peak) (cm^{-2})	3.5×10^{19}	6.0×10^{19}	17.0×10^{19}	2.5×10^{19}
L_x (pc)	90–200	50–140	340–530	60–130
L_y (pc)	1000–2500	100–400	1400–2100	60–130
$n(\text{H I})$ (cm^{-3}).....	0.13–0.06	0.37–0.14	0.16–0.10	0.13–0.06
$M(\text{H I})$ (M_\odot).....	$(0.73\text{--}4.02) \times 10^5$	$(1.86\text{--}13.57) \times 10^4$	$(0.68\text{--}1.60) \times 10^6$	$(0.09\text{--}0.41) \times 10^4$
$M(\text{H I})$ ($M_\odot \text{ yr}^{-1}$).....	$(2.0\text{--}4.7) \times 10^{-3}$	$(16.0\text{--}43.2) \times 10^{-4}$	$(6.4\text{--}9.8) \times 10^{-3}$	$(0.8\text{--}1.6) \times 10^{-4}$
H ⁺ Parameters Assuming that H ⁺ Has Same Path Length as H I				
$I(\text{H}\alpha)$ (R).....	0.07 ± 0.01	<0.15	0.27 ± 0.03	1.1 ± 0.2
$n(\text{H}^+)$ (cm^{-3}).....	0.04–0.03	<0.08–0.05	0.04–0.03	0.20–0.14
$N(\text{H}^+)$ (cm^{-2})	$(1.1\text{--}1.7) \times 10^{19}$	< $(1.3\text{--}2.1) \times 10^{19}$	$(4.4\text{--}5.5) \times 10^{19}$	$(3.8\text{--}5.6) \times 10^{19}$
$M(\text{H}^+)$ (M_\odot).....	$(0.7\text{--}5.9) \times 10^5$	< $(0.7\text{--}8.5) \times 10^4$	$(0.6\text{--}1.9) \times 10^6$	$(0.4\text{--}2.3) \times 10^4$
$M(\text{H}^+)$ ($M_\odot \text{ yr}^{-1}$).....	$(1.9\text{--}6.8) \times 10^{-3}$	< $(6.0\text{--}26.8) \times 10^{-4}$	$(6.7\text{--}12.8) \times 10^{-3}$	$(3.0\text{--}9.2) \times 10^{-4}$
H ⁺ Parameters Assuming that H ⁺ Has Same Volume Density as H I				
L (pc).....	9–51	<2–18	23–55	148–660
$n(\text{H}^+)$ (cm^{-3}).....	0.13–0.06	<0.37–0.14	0.16–0.10	0.13–0.06
$N(\text{H}^+)$ (cm^{-2})	$(0.4\text{--}0.9) \times 10^{19}$	< $(0.3\text{--}0.8) \times 10^{19}$	$(1.2\text{--}1.8) \times 10^{19}$	$(5.9\text{--}12.4) \times 10^{19}$
$M(\text{H}^+)$ (M_\odot).....	$(0.2\text{--}2.9) \times 10^5$	< $(0.2\text{--}3.0) \times 10^4$	$(0.2\text{--}0.6) \times 10^6$	$(0.6\text{--}5.2) \times 10^4$
$M(\text{H}^+)$ ($M_\odot \text{ yr}^{-1}$).....	$(0.6\text{--}3.4) \times 10^{-3}$	< $(1.3\text{--}9.5) \times 10^{-4}$	$(1.7\text{--}4.1) \times 10^{-3}$	$(4.6\text{--}20.5) \times 10^{-4}$

NOTE.—The mass flows are for infall in the case of the Cohen Stream and the Perseus arm IVC, but they represent outflows for Complex GCP and cloud g1.

The resulting column densities are divided by $N(\text{H I})$, giving the values in columns (11) and (12). We note that the error in the Ca II/H I ratio is calculated using the observed error in the Ca II column density combined with an assumed 20% error in $N(\text{H I})$.

Column (13) shows the final result derived from each star: a “U” if the star yields an upper distance limit, an “L” if it gives a lower limit, and an ellipse if the S/N was insufficient to derive a lower limit from a nondetection. The significance of the non-detections is obtained by comparing the expected value (cols. [5] and [6]) with the 3σ upper limits (cols. [7] and [8]). As discussed by Wakker (2001), a safety factor needs to be taken into account for such a comparison, since we do not precisely know the H I column density (see discussion of col. [4] above) and the Ca II/H I ratio in the cloud may in principle vary due to variations in calcium depletion and ionization conditions. We assume a factor of ~ 1.5 for the uncertainty in $N(\text{H I})$ and another factor of ~ 2 for the uncertainty in the Ca abundance. Only if the ratio $\text{EW}(\text{expected})/\sigma(\text{EW})$ is >9 do we conclude that a nondetection sets a lower limit to the cloud’s distance.

4.3. Calculation of Masses and Mass Flow Rates

After we determine the distance to a cloud, we can estimate its location in the Milky Way, its mass, and the mass flow rate associated with it. These numbers are summarized in Table 4.

We estimate the mass in the manner described by Wakker & van Woerden (1991). In summary, we find the values of T_B for each point in the cloud that were given by Hulsbosch & Wakker (1988). Converting T_B to a column density (using the line width) and then multiplying by the area (in cm^{-2}) represented by one

survey grid cell (about 1 deg^2) gives the H I mass in the grid cell. Integrating over the cloud gives the total mass, given an assumed distance, D . This value scales as D^2 .

We use the Hulsbosch & Wakker (1988) data set instead of the newer LAB data from Kalberla et al. (2005) because in the latter we have to select a fixed velocity range, so there can be blending with lower velocity gas. Also, the cloud edges need to be defined by polygons. The older survey clearly separates any particular HVCs from other gas. An improved list could be constructed from the LAB survey, but this has not yet been done. Where possible we do check the Hulsbosch & Wakker (1988) cloud masses against a rough estimate using the LAB survey, and we find that they are similar to about 25%.

To derive the mass flow rate, we need to estimate a vertical velocity for each survey grid cell. Given the observed LSR velocity, we calculate the “deviation velocity” (v_{dev}), the amount by which the cloud’s velocity differs from the maximum velocity that can be understood from a simple model of differential galactic rotation (see Wakker 1991). This simple model consists of a disk with radius 26 kpc with thickness 4 kpc inside the solar circle (8.5 kpc) increasing to 12 kpc at 26 kpc. The rotation curve is flat, except in the inner 0.5 kpc where solid-body rotation is assumed. Using the deviation velocity, we estimate the cloud’s z -velocity (v_z) by assuming that it is purely vertical or by assuming that it is purely radial with respect to the Sun. In the former case, $v_z = v_{\text{dev}}/\sin b$, while in the latter $v_z = v_{\text{dev}} \sin b$. To estimate the mass flow, we take the average of these two possibilities. Dividing the cloud’s distance by the estimated vertical velocity yields the time it will take for the cloud to reach the Galactic plane. Finally, the quotient




FIG. 2.—Map of the column density of the Cohen Stream, integrating the LAB data of Kalberla et al. (2005) between $v_{\text{LSR}} = -130$ and -80 km s^{-1} . Contours are at column densities of 1×10^{19} , 2×10^{19} , and $4 \times 10^{19} \text{ cm}^{-2}$. The locations of the background stars are overlaid. The size (i.e., area) of the symbols is proportional to the distance of the star. Filled circles are for directions where the HVC is detected in absorption. Filled triangles are for significant nondetections. Open circles are for non-detections that do not allow a conclusion concerning the HVC's distance. Crosses are for stars where the interstellar absorption is too blended with stellar lines. The QSO is shown by a five-pointed star instead of a filled circle.

of cloud mass and travel time gives the mass flow associated with the cloud. The resulting number scales as D .

4.4. Ionized Hydrogen

To fully understand the clouds and estimate the mass flows associated with them, we also need to estimate the amount of ionized hydrogen. It turns out that $\text{H}\alpha$ data exist for each of our clouds. There are a number of single pointings for Complex GCP and the Cohen Stream, and there are maps from the Wisconsin $\text{H}\alpha$ Mapper (WHAM) northern sky survey (Haffner et al. 2003) for cloud g1 and the Cohen Stream field IVC. We describe these data when discussing each cloud.

Since we know the cloud distances, we can use the $\text{H}\alpha$ data to estimate the amount of ionized hydrogen in our clouds. There are two ways to do this. In both cases we assume that $N(\text{H}^+)$ is constant, which is justified if the ionization is due to photons from the Galactic disk: hydrogen ionization continues until all Lyman continuum photons have been soaked up. Then, we can first assume that the ionized hydrogen is distributed throughout the cloud, in which case the cloud thickness we use to convert the emission measure to a density is similar to its width on the sky. We can also assume that all the ionized hydrogen sits on the surface of the cloud and the cloud's density is constant; i.e., we use $n(\text{H}^+) = n(\text{H I})$ and calculate the emission length from the emission measure. Both these methods are used to derive the numbers presented in Table 4.

5. RESULTS

5.1. Cohen Stream

5.1.1. Distance

The first cloud we observed is part of the so-called Cohen Stream, discovered by Cohen (1981) (although he only observed the part of the stream that lies to the east of our stars). This HVC is a rather linear stream of gas that runs from $(l, b) \sim (175^\circ, -30^\circ)$ to $(l, b) \sim (140^\circ, -52^\circ)$, with velocities of $v_{\text{LSR}} \sim -110 \text{ km s}^{-1}$. It lies in a part of the sky where the maximum velocity expected from differential galactic rotation is -15 km s^{-1} . A partial map of the cloud is given in Figure 2. On its eastern end ($l > 156^\circ$) it overlaps with a bright HVC at $v_{\text{LSR}} = -280 \text{ km s}^{-1}$, also discovered by Cohen (1981), and named WW 507 by Wakker & van Woerden (1991). It is still unclear whether these two HVCs are related or whether they are a chance coincidence. Unfortunately, the footprint of the SDSS does not overlap the WW 507 cloud, so we do not (yet) know distant stars in its direction. We return to this point in § 5.1.5.

We obtained VLT UVES spectra for nine stars and one QSO in the direction of the Cohen Stream, shown in Figure 3. Seven of these objects lie in the direction of the small core at the end of the stream (cloud WW 516 in Wakker & van Woerden 1991), and three lie projected onto its main body (cloud WW 517). As we discuss below, and as can be seen in Figure 3 and Table 3, we detect high-velocity interstellar Ca II H and K absorption toward




FIG. 3.—Spectra for stars toward the Cohen Stream. Flux units (f.u.) are 10^{-16} ergs cm^{-2} s^{-1} \AA^{-1} . *Left*: Ca II K spectra (histograms) and continuum fits (solid lines); *middle*: Ca II H spectra and continua (note that the left flux scale is valid only for Ca II K); *right*: H I 21 cm spectra. One star is given per row, with the name and distance given in the top left corner of the K and H panels. Distances are preceded by “>” for lower limits and by “<” for upper limits. Distances in brackets show that the detection limit is worse than the expected equivalent width, or there are too many stellar lines. Labels [Ca II (*)] and [He I (*)] show the positions of stellar absorption lines. Vertical lines give the velocities of the H I components. Triangles and numbers in each Ca II panel give the shape of the expected absorption line, the expected equivalent width, and the 3σ detection limit or the detected equivalent width (a “0” means that there is a blended stellar line).

the QSO SDSS J014631.99+133506.3, as well as the K line toward the 11.2 ± 1.0 kpc distant RR Lyrae star SDSS J015133.91+141105.2. The most distant star toward which we do not detect the HVC at a significant level is the BHB star 2MASS J014936.48+143914.63 at 4.6 ± 0.8 kpc. Ca II is also not detected toward SDSS 014843.61+130411.5 ($D = 5.0 \pm 1.2$ kpc) and SDSS 020033.50+141154.0 ($D = 8.7 \pm 1.3$ kpc). The first of these is not significant, although just barely so, while for the latter the ratio of expected to

observed equivalent width is only 1. Taking into account the errors on the star’s distances, these results imply a distance bracket of 5.0 to 11.7 kpc for the Cohen Stream. If we relax the criterion for a significant nondetection, we could improve the lower limit to 5.6 kpc.

5.1.2. Discussion of Stars

1. SDSS J015133.91+141105.2 is the star that sets the upper limit on the distance. The HVC K line is clearly seen in absorption

in its spectrum, with an equivalent width of $17 \pm 3 \text{ m}\text{\AA}$. This low-metallicity ($[\text{Fe}/\text{H}] = -1.73 \pm 0.3$) AV star shows relatively few stellar lines, and with its velocity of -176 km s^{-1} , the stellar Ca II and Fe I $\lambda 3931.409$ lines are shifted well away from the HVC. The expected equivalent width of the H line is $10 \text{ m}\text{\AA}$, far below the $34 \text{ m}\text{\AA}$ detection limit. This detection limit is rather high because the stellar H ϵ line substantially reduces the star's flux at the velocity of the HVC ($13.2 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ at Ca II K, $3.5 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ at Ca II H). Thus, we cannot use the H line to confirm the K line detection. However, we have seven exposures of this star, taken on August 10, September 7 (two exposures), and September 23 (four exposures 1 hr apart). These show the stellar velocity varying over time: $v(\star) = -194 \pm 3 \text{ km s}^{-1}$ on August 10, -138 km s^{-1} on September 7, and $-178, -193, -193,$ and -190 km s^{-1} on September 23. From these spectra it is clear that the feature at $v_{\text{LSR}} = -104 \text{ km s}^{-1}$ is interstellar, as it stays at the same position, whereas the stellar lines vary. In order to have a clean continuum against which the HVC absorption can be measured, we combined only the first and last exposure, which have the cleanest separation between stellar and interstellar absorption.

2. Toward the QSO SDSS J014631.99+133506.3 the HVC's K and H equivalent widths are 17 ± 3 and $11 \pm 3 \text{ m}\text{\AA}$, respectively. The Ca II/H I ratio implied by the K line is about half the value found toward the star SDSS J15133.91+141105.2. They differ at the 2σ level, but this can easily be explained by the large uncertainty in the value of $N(\text{H I})$ toward the two targets. On the other hand, the equivalent widths of the K and H lines seen toward the QSO are in the expected ratio of about 2, giving confidence in the detections.

3. Using the method described in § 2, we predict K and H equivalent widths of 17 and $8 \text{ m}\text{\AA}$ toward the $30.4 \pm 3.9 \text{ kpc}$ distant BHB star SDSS J015735.65+135254.2. In its spectrum we find a possible HVC K absorption of $10 \pm 3 \text{ m}\text{\AA}$. Since it is just a 3σ feature, we do not claim this as a clear detection of the HVC, but the feature is compatible with the expectations.

4. Toward the star 2MASS J01449.36.48+143914.6 ($D = 4.6 \pm 0.8 \text{ kpc}$) the ratio of expected Ca II K equivalent width ($21 \text{ m}\text{\AA}$) to the observed error ($2.9 \text{ m}\text{\AA}$) is 7.2, which implies that this star sets a lower distance limit to the distance of the HVC.

5. The HVC is also not seen in absorption toward the other six stars with $D < 10 \text{ kpc}$. Toward most of these stars the H I column density is low, and only the nondetections toward the four nearest stars are significant.

5.1.3. Ca II/H I Ratio

The Ca II/H I ratio that we find for the Cohen Stream is on the low end of those found for HVCs: $(11.7 \pm 2.9) \times 10^{-9}$ toward SDSS J015133.91+141105.2 and $(6.8 \pm 2.0) \times 10^{-9}$ toward SDSS J014631.99+133506.3. The Ca II/H I ratios predicted by the Wakker & Mathis (2000) correlation are 27×10^{-9} and 17×10^{-9} , respectively. The ratio of observed to expected values thus is 0.42 ± 0.05 ; i.e., the Cohen Stream appears to be relatively underabundant by a factor of ~ 2.4 in gaseous ionized calcium. However, until we have a measurement of the metallicity of the Cohen Stream, it will remain unclear whether this underabundance is due to low metallicity, unusually high calcium ionization, or high depletion of calcium onto dust, or even small-scale structure in the H I, which affects our estimate of $N(\text{H I})$.

5.1.4. Location in the Milky Way

Previous assessments of the distance of the Cohen Stream were summarized by Wakker (2001): Kemp et al. (1994) and Tamanaha (1996) observed Mg II, Ca II, and Na I in bright stars in the direc-

tion of the Cohen Stream, at distances up to 300 pc , but no absorption was found. Based on a morphological study of the high-, intermediate-, and low-velocity gas, Tamanaha (1995) argued that the Cohen Stream is the remnant of a collision between the WW 507 cloud and low-velocity gas, which would place it within about 1 kpc from the disk. Our distance bracket excludes such a simple interaction model, as the cloud is clearly more distant than 9.2 kpc .

It is now clear that this cloud is rather far below the Galactic plane, having z between 3.7 and 8.6 kpc (see Table 4). It is located at similar longitude, but on the opposite side of the plane and further inward than the Galactic warp (which is at galactocentric radii between about 17 and 22 kpc ; Burton & de Lintell Hekkert 1986).

5.1.5. Ionized Hydrogen

Weiner et al. (2001) reported H α emission from two directions in the anticenter region. One of these is at $(l, b) = (164^\circ, -46^\circ)$ and samples the Cohen Stream (B. J. Weiner 2007, private communication). The H α intensity $[I(\text{H}\alpha)]$ is $0.07 \pm 0.01 \text{ R}$. [One Rayleigh (R) is $10^6/4\pi \text{ photons cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1}$; the emission measure (EM = $\int n_e^2 L$) is related to this as $\text{EM} = 2.75T_4^{0.924}I(\text{H}\alpha)$, where T_4 is the gas temperature in units of 10^4 K .]

The H $^+$ column density we derive from the observed H α emission is on the order of 10^{19} cm^{-2} , independent of most assumptions. The H $^+$ mass is found to be similar to the H I mass, which is a few times $10^5 M_\odot$. If we assume that the H $^+$ is located in a layer around the neutral cloud, the implied skin thickness is 10% to 25% of the total cloud thickness.

We noted earlier that H α emission was also detected from the higher velocity cloud near the Cohen Stream (Kutuyev & Reynolds 1989; Weiner et al. 2001), with an intensity of $\sim 0.08 \text{ R}$. This cloud is centered around $l = 168^\circ, b = -46^\circ, v_{\text{LSR}} = -280 \text{ km s}^{-1}$. Using the Bland-Hawthorn & Maloney (1999) model of the Galactic ionizing radiation field, the H α brightness of this cloud would imply a distance on the order of 15 kpc . If such an interpretation for the H α emission can be maintained, this would suggest that this cloud and the Cohen Stream are close together in space, even though their velocities differ by 170 km s^{-1} .

5.2. High- z Perseus Arm IVC

5.2.1. Distance

In the sight lines to the Cohen Stream an IVC is also visible in the H I spectra, with velocities of roughly -40 to -60 km s^{-1} . This IVC is part of an extended region of intermediate-velocity gas, a map of which can be found in Wakker (2001, their Fig. 18), who named this gas "IV-south." Figure 4 shows the H I with v_{LSR} between -80 and -30 km s^{-1} . In the general vicinity of this IVC differential galactic rotation contributes substantially to the observed radial velocity (up to -15 km s^{-1} at $b = -50^\circ$, up to -45 km s^{-1} at $b = -25^\circ$). Therefore, we present in Figure 5 the H I column density for deviation velocities between -60 and -30 km s^{-1} , i.e., gas parcels whose velocities are 30 to 60 km s^{-1} more negative than the maximum radial velocity expected from differential galactic rotation. This clearly shows the IVC centered near $(l, b) = (145^\circ, -42^\circ)$. The stellar spectra for targets projected onto the IVC can be seen in Figure 3. The stream running from $(l, b) \sim (130^\circ, -60^\circ)$ to $(l, b) \sim (100^\circ, -47^\circ)$ was termed the "Pegasus-Pisces Arch" or PP Arch by Wakker (2001), and its distance is $< 2.7 \text{ kpc}$.

Intermediate-velocity absorption is clearly detected toward the QSO and in the spectra of all stars more distant than 2.1 kpc , but not toward PG 0142+148 ($D = 1.2 \pm 0.5 \text{ kpc}$), where the ratio $\text{EW}(\text{expected})/\sigma(\text{EW})$ is 12.7 . Taking into account the errors on the stellar distances, the implied distance bracket is 1.0 to 2.7 kpc .




FIG. 4.—Map of the column density of the IVC in the Cohen Stream field, gas probably associated with the high- z extension of the Perseus arm, integrating the LAB data of Kalberla et al. (2005) between $v_{\text{LSR}} = -80$ and -30 km s^{-1} . Contours are at column densities of 1×10^{19} , 2×10^{19} , and 4×10^{19} cm^{-2} . The locations of the background stars are overlaid. Symbols are as for Fig. 2.




FIG. 5.—Map of the column density of the intermediate-velocity gas in a large region to the south of the Perseus arm. This map represents the HI column density in the LAB data of Kalberla et al. (2005) integrated between deviation velocities $v_{\text{dev}} = -60$ and -30 km s^{-1} . Contours are at column densities of 1×10^{19} , 2×10^{19} , and 4×10^{19} cm^{-2} . The box outlines the region shown in Figs. 2 and 4.

5.2.2. Ca II/H I Ratio

For this IVC the observed Ca II/H I ratios range from $(10 \pm 3) \times 10^{-9}$ toward 2MASS J014936.48+143914.6 to 30 ± 9 toward SDSS J015735.65+135254.2. Much of this range might be due to the uncertainty in the H I column densities toward the stars. The ratio of observed to predicted Ca II/H I ratio is on average 0.62 ± 0.11 , ranging from 0.41 to 0.75. Thus, like the Cohen Stream, this gas also seems to be deficient in ionized calcium compared to the ratio $N(\text{Ca II})/N(\text{H I})$ for HVCs in general, but we do not have enough information to determine the cause of this.

5.2.3. Location in the Milky Way

The IVC lies below the Perseus arm, the next spiral arm out from the Local arm. In the Galactic plane, the Perseus spiral arm has velocities of about -50 km s^{-1} . The distance to the Perseus arm is about $2.5 \pm 0.5 \text{ kpc}$ (see Reynolds et al. 1995 and references therein). Haffner et al. (1999) studied the H α emission from the high- z extension of the Perseus arm and find that it has a scale height of about 1 kpc. Associated gas at $b = -45^\circ$ would be at a distance of $3.6 \pm 0.7 \text{ kpc}$ and lie $2.5 \pm 0.5 \text{ kpc}$ below the Galactic plane. A z -height of 1 kpc implies a distance of 1.6 kpc, in which case the cloud would be located in the interarm region. Gas near the upper distance limit of 2.7 kpc would have $z = 1.9 \text{ kpc}$. So, the IVC we see is likely associated with the front side of the high- z extension of the Perseus arm.

For this direction, the LSR velocity predicted from differential galactic rotation is roughly -25 km s^{-1} , suggesting that the IVC has a peculiar velocity toward the plane of between 15 and 35 km s^{-1} . The derived distance bracket is not very precise, but it is clearly compatible with the idea that this IVC is associated with the Perseus arm. This could be confirmed by obtaining spectra with high S/N of additional stars in the 1 to 3 kpc range.

To estimate the cloud's H I mass, we integrated the column densities between $(l, b) \sim (139^\circ, -47^\circ)$ and $(l, b) \sim (155^\circ, -40^\circ)$. The cloud's total mass is found to be in the range $(0.2\text{--}1.4) \times 10^4 M_\odot$. We note that this is of the same order of magnitude as the ‘‘Intermediate-Velocity Arch’’ (IV Arch), the largest northern IVC (see Kuntz & Danly 1996; Wakker 2001, 2004), which also is moving toward the Galactic plane with a velocity of $\sim 40 \text{ km s}^{-1}$. It is also similar to the mass of the ‘‘Low-Latitude Intermediate-Velocity Arch’’ (LLIV Arch), an IVC that is 0.9–1.8 kpc distant with a mass of $1.5 \times 10^5 M_\odot$. The latter IVC appears to be in the interarm region. The IVC we detected in absorption may be a cloud similar to the IV and LLIV Arches, but associated with the Perseus arm, rather than with the Local spiral arm.

5.2.4. Ionized Hydrogen

In the WHAM H α survey (Haffner et al. 2003), there is no detectable H α emission from the area covered by this cloud, which corresponds to a 3σ upper limit of $I(\text{H}\alpha) < 0.15 \text{ R}$, or an emission measure $< 0.07 \text{ cm}^{-6} \text{ pc}$. The implied H $^+$ column density limit of $\sim 10^{19} \text{ cm}^{-2}$ is compatible with the values found for the other clouds, so the nondetection does not imply a lack of ionized hydrogen for this object.

The nondetection does imply that for this cloud the mass of ionized gas is smaller than the neutral hydrogen mass. If the ionized gas is on a skin, this skin is relatively thin, being at most 10% of the cloud's thickness.

5.3. HVC Complex GCP or HVC 40–15+100

5.3.1. Distance

The second HVC in our sample lies near $l = 40^\circ$, $b = -15^\circ$ and has $v_{\text{LSR}} \sim 90 \text{ km s}^{-1}$. It has been referred to as the ‘‘Smith

Cloud,’’ after Smith (1963), who discovered it, or as HVC 40–15+100, or as Complex GCP (Wakker & van Woerden 1991). Wakker (2001) could only set a lower distance limit of 0.3 kpc, from the nondetection of Ca II absorption toward some nearby stars.

We observed eight stars projected on Complex GCP (see Fig. 6), six of which were found from 2MASS; the other two are RR Lyrae stars from the Kukarkin et al. (1970) catalog. The Ca II absorption and H I emission spectra are shown in Figure 7. We detect the HVC in the spectrum of V1084 Aql, located at $14.5 \pm 1.3 \text{ kpc}$, but not toward V1172 Aql at $10.5 \pm 1.3 \text{ kpc}$, yielding a distance bracket of 9.8 to 15.1 kpc.

5.3.2. Discussion of Stars

1. Figure 7 shows the HVC absorption in the V1084 Aql K line, which has an equivalent width of $51 \pm 3 \text{ m}\text{\AA}$. We caught this low-metallicity ($[\text{Fe}/\text{H}] = -1.22 \pm 0.25$) RR Lyrae star at a temperature of about 7000 K, and the spectrum shows relatively few stellar lines. The Ca II H line detection is less clear ($37 \pm 10 \text{ m}\text{\AA}$), not only because it is confused by the stellar H ϵ line, but also because the H ϵ lines makes the flux much lower at the HVC's velocity ($8.5 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ at K, $1.5 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1} \text{ \AA}^{-1}$ at H).

2. The HVC is absent in the spectrum of V1172 Aql ($D = 10.5 \pm 1.4 \text{ kpc}$), and the ratio expected/error is 15.5 for Ca II K and 7.1 for Ca II H. This star thus yields a lower limit for the HVC's distance. We note that the spectrum of V1172 Aql shows weak ($\sim 25 \text{ m}\text{\AA}$) wiggles at the velocity of the HVC in both the K and the H line, but these features do not match up as interstellar Ca II: they are not at exactly the same velocity, and they are not in the ratio 2 : 1.

3. One other star has a large distance (2MASS J195922.75+000300.0 at $9.0 \pm 3.7 \text{ kpc}$), but it has low UV flux as it is a cool star ($T_{\text{eff}} = 5750 \text{ K}$), so the nondetection is not significant.

4. The remaining four stars all have distances less than 2 kpc. The nondetections of interstellar Ca II K toward 2MASS J195927.29+000822.3 and 2MASS J195912.00–002645.3 are significant, but for 2MASS J195741.61–004009.7 and 2MASS J195925.49–000519.1 this is not the case. Considering the distance bracket we have set, these nondetections are not unexpected since these stars are relatively nearby.

5. Finally, one star in our sample is at about the same distance as V1084 Aql: 2MASS J195823.36–002719.0, at $D = 13.1 \pm 3.2 \text{ kpc}$. The 2MASS catalog gives a J magnitude of 15.1 for this star, but our photometry (R. Wilhelm et al. 2008, in preparation) yields $V = 16.9$. Since we expected the star to be bright (and originally estimated a distance of 5.8 kpc), we only requested a short exposure time (7200 s). Even less data were actually obtained (2665 s), so the spectrum has an S/N of only 5 at the location of the interstellar K line. Therefore, the nondetection is not significant (61 m \AA expected, 3σ upper limit 67 m \AA).

5.3.3. Ca II/H I Ratio

Weighing the K line twice as much as the H line, the average Ca II/H I ratio in Complex GCP that we find from the K and H line detected toward V1084 Aql is $(6.2 \pm 1.7) \times 10^{-9}$. If we estimate EW(K) from the Wakker & Mathis (2000) relation between $\log R(\text{Ca II})$ and $\log N(\text{H I})$, we predict $R(\text{Ca II}) \sim 6 \times 10^{-9}$ for Complex GCP. Thus, the Ca II/H I ratio is exactly what is expected.

5.3.4. Location in the Milky Way

We place Complex GCP at a distance of 9.8 to 15.1 kpc, which puts it at a galactocentric radius between 6.2 and 9.7 kpc, and




FIG. 6.—Map of the column density of Complex GCP, integrating the LAB data of Kalberla et al. (2005) between $v_{\text{LSR}} = +60$ and $+130$ km s $^{-1}$. Contours are at $N(\text{H I}) = (1, 2, 4, 10, 20, 30) \times 10^{19}$ cm $^{-2}$. Symbols are as for Fig. 2.

2.5 to 3.9 kpc below the plane. Its $2^\circ \times 8^\circ$ size on the sky implies a linear size of about 0.5×1.8 kpc. The H I mass is about $10^6 M_\odot$.

Bland-Hawthorn et al. (1998) suggested that this HVC is related to the Sagittarius dwarf, at a distance of 26 ± 4 kpc ($R = 20$ kpc, $z = -7$ kpc). We find that instead of lying on the outskirts of the Milky Way, this HVC is near the solar circle, only halfway to the Sagittarius dwarf. Only a detailed model of the dwarf galaxy's orbit can tell whether the HVC could be related after all.

In the direction of Complex GCP differential galactic rotation can explain a velocity of up to $+75$ km s $^{-1}$, which occurs at the tangent point ($D = 7$ kpc, at galactocentric radius 5.5 kpc, 2.0 kpc below the plane). At distances of 9.8 and 15.1 kpc, gas in differential rotation has an LSR velocity of $+50$ and -17 km s $^{-1}$, respectively. It is therefore clear that Complex GCP is below the Galactic plane and moving ahead of the gas in the plane with a velocity of 40 to 105 km s $^{-1}$. A. J. Heroux et al. (2008, in preparation) mapped this HVC using the new Green Bank Telescope. They are working on a model of its orbit, including drag forces to explain the stretching that is obvious in the sky map (Fig. 6).

5.3.5. Ionized Hydrogen

Bland-Hawthorn et al. (1998) based their suggested association with the Sagittarius dwarf on the detected H α emission from this HVC, with intensities $0.24 R$ at $(l, b) = (40^\circ, -15^\circ)$ and $0.30 R$ at $(l, b) = (41^\circ, -14^\circ)$. If we assume that the cloud's thickness is comparable to the width of its short dimension (0.3–0.5 kpc), these emission measures imply an electron density n_e of

$0.04\text{--}0.03$ cm $^{-3}$, about 25% of the implied H I density of $0.16\text{--}0.10$ cm $^{-3}$. In that case the H $^+$ mass is similar to the cloud's H I mass. If, on the other hand, we assume that the H $^+$ occurs around the outside of the cloud, the ionized mass is about one-third of the neutral mass, while the skin is about 10% of the thickness of the cloud. In either case it is clear that this cloud must have a substantial ionized component. Now that we know its location, a full H α map would clearly help understand the Galactic ionizing radiation field.

5.4. Cloud g1

5.4.1. Distance

The last cloud in our sample is an IVC near $(l, b) = (67^\circ, -27^\circ)$, at $v_{\text{LSR}} \sim +70$ km s $^{-1}$ (Fig. 8). It is one of a set of faint positive-velocity IVCs in the region $l = 30^\circ\text{--}70^\circ$, $b = -45^\circ$ to -20° that Wakker (2001) termed “Complex gp.” For the present project we have designated some of the cores in this complex as g1 through g5, with the IVC toward which we observed stars termed “cloud g1.” It is, a priori, unclear whether these IVCs are related to Complex GCP, which lies near the IVCs, although at much higher positive velocity.

Many stars projected onto cloud g1 have previously been observed. It was detected in Ca II and Na I absorption toward several stars in M15 ($D = 10$ kpc; Cohen 1979; Songaila & York 1980; Kennedy et al. 1998; Lehner et al. 1999). Intermediate-velocity absorption at the same velocity as cloud g1 was also seen toward the star HD 203664 ($D = 4.3 \pm 0.4$ kpc; Albert et al. 1993; Little




FIG. 7.—Same as Fig. 3, except for Complex GCP.

et al. 1994; Ryans et al. 1996), which lies 3° west of the main cloud core (see Fig. 8). Absorption is not detected toward HD 203699 ($D = 0.8 \pm 0.1$ kpc; Kennedy et al. 1998), nor toward several A and F stars at distances between 0.2 and 1.4 kpc (Smoker et al. 2001). The nondetections in the latter paper are usually not significant by our criteria, except maybe for their star 7, an A2 V star at a distance of 1.4 kpc. Together, these detections and nondetections bracketed the distance of g1 as 0.7 (maybe 1.4) to 4.5 kpc, with the caveat that these stars do not lie near the brighter part of the cloud but probe the diffuse distribution of gas with similar velocities.

Smoker et al. (2002) made a detailed study of the part of the cloud that lies in front of M15. They combined high angular resolution ($1' \times 2'$) H I (Westerbork+Arcibo) data with WHAM H α profiles and an *IRAS* map of the region. They found that the H I has structure on scales of $5'$ ($1.5D$ [kpc] pc) and that the gas is warm (~ 500 K).

We observed seven stars projected onto the main core of cloud g1, with distances ranging from 1.2 to 8.2 kpc. Since these stars are relatively nearby, the spectra have relatively high S/Ns. Absorption is seen toward three stars, 2MASS J213327.04+133026.7 ($D = 8.2 \pm 0.9$ kpc), FW Peg ($D = 3.8 \pm 0.8$ kpc),

and BS 17578–0016 ($D = 3.6 \pm 0.5$), while significant non-detections are obtained toward the other four stars (ratios of expected EW to upper limit of 25 to 60 for Ca II K). The most distant of these is PG 2134+125 at 2.2 ± 0.8 kpc. Using the 1σ range allowed by the stellar distances, the distance of cloud g1 is then bracketed as 1.8 to 3.8 kpc. This is a range of 2 kpc, tightening the previous 3.8 kpc wide bracket. Unfortunately, the current distance bracket still means that the derived cloud parameters will have an uncertainty of a factor of 2 to 4 (depending on how they scale with distance).

5.4.2. Discussion of Stars

1. As Figure 9 shows, the four nondetections toward BS 17578–0015, 2MASS J213520.04+133045.0, 2MASS J213451.58+134017.5, and PG 2134+125 are very clear. The spectra show no stellar lines near the HVC velocity, and the stellar Ca II K, H, and He I lines are not interfering.

2. The detection toward FW Peg is simple and certain: the Ca II K and H absorption lines have equivalent width ratio 2, to within the errors. The stellar velocity of -225 km s $^{-1}$ shifts the stellar Ca II K nicely out of the way. Further, we have two spectra of this star (taken 16 days apart) in which the stellar lines clearly




FIG. 8.—Map of the column density of the g1 cloud, integrating the LAB data of Kalberla et al. (2005) between $v_{\text{LSR}} = +55$ and $+90$ km s $^{-1}$. Contours are at 1×10^{19} , 2×10^{19} , and 4×10^{19} cm $^{-2}$. Symbols are as for Fig. 2.

shift in velocity, but the lines interpreted as interstellar do not. A few (unidentified) stellar lines can be seen toward more positive velocities than the IVC absorption, but they do not confuse the issue.

3. The detection toward 2MASS J213327.04+133026.7 is not as clean as the one toward FW Peg, but the line is obviously interstellar, since the spectrum shows no stellar lines other than Ca II K, H, and H ϵ . However, the H ϵ line is very broad and deep, lowering the flux near the velocity of the interstellar absorption and making the detection of Ca II H more noisy.

4. The most problematic star is BS 17578–0016. Both the H I 21 cm profile and the Ca II absorption lines show two narrow components, separated at the velocity where the H I in g1 usually peaks. This actually gives confidence in the interpretation of the absorption as associated with g1, but it makes the measurements more difficult. In Table 3 we list the absorption as two separate components.

5.4.3. Ca II/H I Ratio

As summarized by Wakker (2001), Lehner et al. (1999) detected Ca II in cloud g1 toward 13 stars in M15. Using values of $N(\text{H I})$ between 42×10^{18} and 54×10^{18} cm $^{-2}$ for these stars gives Ca II/H I ratios between 14×10^{-9} and 79×10^{-9} , averaging to 35×10^{-9} , although individual measurements have large errors. Toward HD 203664 Ryans et al. (1996) derived a Ca II/H I ratio of $(450 \pm 40) \times 10^{-9}$ [for $N(\text{H I}) = 2.2 \times 10^{18}$ cm $^{-2}$]. For the three stars toward which we detect cloud g1 we find values of $(106 \pm 22) \times 10^{-9}$ (FW Peg), $(63 \pm 16) \times 10^{-9}$ (2MASS J213327.04+133026.7), $(109 \pm 28) \times 10^{-9}$ (BS 17578–0016, stronger H I

component), and $(510 \pm 110) \times 10^{-9}$ (BS 17578–0016, weaker H I component). Using the Wakker & Mathis (2000) relation, we would predict Ca II/H I ratios of 11×10^{-9} for the stars in M15, 140×10^{-9} toward HD 203664, and 28×10^{-9} , 35×10^{-9} , 103×10^{-9} , and 225×10^{-9} for our detections. The ratio of predicted to observed Ca II/H I ratio thus varies from 1.1 for one of the BS 17578–0016 components to 3.8 for FW Peg. However, the H I column densities are comparatively uncertain, since the cloud is known to have structure on much smaller scales than the 36' Dwingeloo beam (Smoker et al. 2002). So, a proper measurement of the Ca II/H I ratios in cloud g1 will require H I observations with much higher resolution toward each of the stellar targets. Nevertheless, it appears that the Ca II/H I ratio is higher than average in cloud g1. Further, the apparently extremely high Ca II/H I ratios for the weak components toward BS 17578–0016 are in line with the values expected from the Wakker & Mathis (2000) relation.

5.4.4. Location in the Milky Way

The distance bracket of 1.8–3.8 kpc for cloud g1 places it at a galactocentric radius of ~ 7.9 kpc and $z = -0.8$ to -1.7 kpc, moving with $v = 55$ km s $^{-1}$ relative to corotating Galactic gas around it. The ranges in the implied galactocentric radius and peculiar velocity are small because g1 lies near the tangent point. Its height below the plane is typical for an IVC. However, unlike the large IV Arch in the northern sky, this gas is moving away from the plane with an estimated velocity of about 25 km s $^{-1}$.

The area of the core of the cloud is only about $2^\circ \times 2^\circ$, which translates to a size of $(60\text{--}130)^2$ pc. The H I mass is only about




FIG. 9.—Same as Fig. 3, except for cloud g1.

1000–4000 M_{\odot} . The neutral part of this cloud is thus clearly fairly small.

5.4.5. Ionized Hydrogen

Smoker et al. (2002) reported $H\alpha$ emission from cloud g1 toward three directions that lie projected onto the areas with the brightest $H\text{ I}$ emission. These detections were taken from the WHAM northern sky survey (Haffner et al. 2003). The $H\alpha$ intensity does not vary much (the detections are 1.30 ± 0.04 , 0.95 ± 0.04 , and 0.91 ± 0.03 R), whereas the $H\text{ I}$ column densities have a range of a factor of 3 for these directions. This result supports the contention that for many HVCs/IVCs $I(H\alpha)$ will be relatively constant across the face of the cloud, and not correlated with $N(H\text{ I})$. With our distance bracket for this cloud, the bright $H\alpha$ intensities imply either a relatively large H^+ volume density ($\sim 0.17\text{ cm}^{-3}$) or a large ionized path length (several hundred parsecs). With either of our assumptions, $M(H^+)$ is much larger than $M(H\text{ I})$, by a factor of 5 to 10. It is clear that what we call cloud g1 is a small neutral condensation in a much larger region of ionized gas.

5.5. Low-Velocity Gas

Low-velocity Ca II absorption is detected toward all stars, although in a few cases stellar lines make it impossible to measure the equivalent widths. It is likely that the low-velocity ab-

sorption is a mixture of several nearby clouds in the line of sight, but we have no way of separating these.

The Ca II/H I ratios measured for the low-velocity gas are similar for the different fields, averaging to $(4.4 \pm 1.6) \times 10^{-9}$ for stars projected on the Cohen Stream, $(5.3 \pm 1.9) \times 10^{-9}$ for the Complex GCP stars, and $(5.7 \pm 2.0) \times 10^{-9}$ for stars in the cloud g1 field. Applying the Wakker & Mathis (2000) relation, these values are only a factor of 2–4 higher than expected. This is remarkable, since that relation was determined using HVC and IVC detections only, and in principle it might not be applicable to low-velocity absorption components since these are probably a mixture of several clouds with different column densities. Also, the column densities of the low-velocity gas are a factor of 10 to 100 larger than the column densities in the HVCs.

In the direction to Complex GCP there is an absorption component at $v_{\text{LSR}} \sim 30\text{ km s}^{-1}$ that is detected in every sight line. This component is well separated in the Ca II absorption, but not in the $H\text{ I}$ profile. To estimate $N(H\text{ I})$, we therefore integrated the $H\text{ I}$ over the same velocity range as the Ca II absorption. The measured Ca II/H I ratio is rather uncertain because the associated $H\text{ I}$ column density is very difficult to measure. Using a flat rotation curve, this gas would have a kinematical distance of 2.0 kpc ($R = 7.1\text{ kpc}$, $z = -0.5\text{ kpc}$). This is consistent with the detection toward the star at $1.8 \pm 0.3\text{ kpc}$, but the detections toward the star that are 0.7 ± 0.6 and $1.1 \pm 0.5\text{ kpc}$ distant suggest that

the gas is much nearer than indicated by the simple kinematical model. At a distance of 0.7 kpc, the expected radial velocity is only $\sim 8 \text{ km s}^{-1}$, so this gas deviates by about 20 km s^{-1} from its expected velocity.

6. DISCUSSION

We now present a short discussion summarizing the implications of our results as they pertain to the mass inflow rate of external gas, to the study of the escape of ionizing radiation from the Milky Way, and to the understanding of tidal streams around the Milky Way. We do not give a full discussion of these items: that will be saved for a future paper, after we have determined more HVC and IVC distances.

One of the more important processes that HVCs trace is the infall of low-metallicity material onto the Milky Way. Wakker et al. (1999) found that HVC Complex C (which has a metallicity 0.15 times solar; Wakker et al. 1999; Richter et al. 2001b; Tripp et al. 2003; Sembach et al. 2004) contributes a mass inflow rate of about $0.1\text{--}0.2 M_{\odot} \text{ yr}^{-1}$, assuming a distance of 10 kpc. A similar calculation for HVC Complex A shows that it contributes about $0.05 M_{\odot} \text{ yr}^{-1}$. Although we do not know the metallicity of the Cohen Stream, its location and velocity lead us to predict that it is a low-metallicity newcomer to the Milky Way. As Table 4 shows, it contributes about $0.01 M_{\odot} \text{ yr}^{-1}$.

There are a number of other HVCs that may represent low-metallicity infalling gas. However, we do not have direct data to prove this. These are the HVCs in Complexes H, ACHV, ACVHV, GN, and P (for the definition see Wakker & van Woerden 1991). If all of these objects were at a distance of 10 kpc, they would represent an inflow rate of $0.4 M_{\odot} \text{ yr}^{-1}$. The Cohen Stream is one of the larger, but not one of the brighter, HVCs in Complex ACHV. If all clouds in this complex are 10 kpc distant, the complex as a whole corresponds to a mass flow rate of $0.1 M_{\odot} \text{ yr}^{-1}$.

Putman et al. (2004) proposed that most, if not all, of the HVCs in the anticenter region are in the orbit of the Sagittarius dwarf, and they suggested that this gas was stripped from that galaxy 200 to 300 Myr ago as it interacted with the gas at the far outskirts of the Galaxy. This would imply a distance on the order of 40 kpc for the anticenter HVCs. We cannot directly exclude that the higher velocity HVC in the region ($v_{\text{LSR}} \sim -280 \text{ km s}^{-1}$) is this distant, but clearly the Cohen Stream is much closer. Moreover, the intensity of the $\text{H}\alpha$ emission from the -280 km s^{-1} cloud is similar to that of the Cohen Stream, which argues that their distances are similar.

We find that the IVC near $(l, b) = (140^{\circ}, -40^{\circ})$ could well be a cloud associated with the Perseus arm. We do not have an accurate distance, but even so we find that its properties are comparable to those of the two large northern IVCs, the IV Arch and the LLIV Arch. Both of these have near-solar metallicity, a mass on the order of $10^5 M_{\odot}$, and each represent an inflow of about $5 \times 10^{-3} M_{\odot} \text{ yr}^{-1}$ (Wakker 2004). Thus, if we were living in the Perseus arm, this IVC would show up as a fairly large solar metallicity infalling cloud near the Galactic south pole.

The g1 cloud has properties just like what is expected for gas taking part in the outflow phase of a Galactic fountain (see Houck & Bregman 1990). As the hot gas flows out, it starts condensing a few kiloparsecs up, before it reaches its maximum height (cloud g1 is at $z \sim 1 \text{ kpc}$, moving out with 25 km s^{-1}). The process of ejecting the gas destroys much of the dust, leading to relatively high gaseous abundances (cloud g1 has a larger than average $\text{Ca II}/\text{H I}$ ratio). But most of the gas is still ionized [$M(\text{H}^+) \sim (5\text{--}10)M(\text{H I})$ for cloud g1]. The only thing we still have not measured directly is the metallicity of cloud g1, which will require spectra in the 1000–1300 Å spectra range for UV-bright stars projected onto it. We expect the metallicity to be solar.

Finally, we note that some of the conclusions above can be confirmed by measuring the cloud metallicities. This can be done by determining the abundance of O I, S II, and other ions in spectra of UV-bright targets located behind our clouds. A check of the *Far Ultraviolet Spectroscopic Explorer (FUSE)* and *Hubble Space Telescope (HST)* archives shows that no useful data exist at present. Targets behind the clouds have been observed, but their spectra have too low resolution, there are too many stellar lines, or they are too faint to provide enough background flux.

7. CONCLUSIONS

We report on VLT observations of 24 stars and one QSO projected on four HVCs. Using these data, we derive distances to these objects. We combine H I and $\text{H}\alpha$ data to estimate the total cloud (neutral and ionized) gas masses. Our conclusions are summarized as follows:

1. We derive a distance bracket of 5.0 to 11.7 kpc for Cohen Stream, a cloud that is the most negative latitude large cloud in the anticenter complex of HVCs. This places the cloud between 3.7 and 8.6 kpc below the Galactic plane. The stream runs from $(l, b) \sim (175^{\circ}, -30^{\circ})$ to $(l, b) \sim (140^{\circ}, -52^{\circ})$ and has velocities of $v_{\text{LSR}} \sim -110 \text{ km s}^{-1}$. Ca II K and H absorption is detected toward the QSO SDSS J014631.99+133506.3 with equivalent widths of 17 ± 3 and $11 \pm 3 \text{ m}\text{\AA}$, respectively. We also find a $17 \pm 3 \text{ m}\text{\AA}$ Ca II K absorption in the spectrum of the $11.2 \pm 1.0 \text{ kpc}$ distant RR Lyrae star SDSS J015133.91+141105.2. The lower limit is derived from the significant nondetection of high-velocity Ca II toward the BHB star 2MASS J014936.48+143914.6 ($21 \text{ m}\text{\AA}$ line expected, 3σ upper limit $9 \text{ m}\text{\AA}$).

2. The distance bracket for the Cohen Stream implies a total neutral hydrogen mass of $(2.5\text{--}3.9) \times 10^5 M_{\odot}$ for the cloud and an associated mass inflow rate on the order of $0.004 M_{\odot} \text{ yr}^{-1}$. $\text{H}\alpha$ emission from the Cohen Stream has been detected at one position, and its intensity suggests an amount of ionized gas that is comparable to the amount of neutral gas. The complex of HVCs that the Cohen Stream is a part of would represent a total inflow of $0.1 M_{\odot} \text{ yr}^{-1}$ if all clouds in the complex are at similar distances. A direct measurement of the metallicity of this complex is still lacking, so the interpretation of the complex as infalling low-metallicity gas is not confirmed.

3. We find a rough distance bracket of 1.0 to 2.7 kpc for intermediate-velocity gas in the direction of the Cohen Stream. This suggests that this gas is part of a high- z extension of the Perseus arm, similar to the more well-studied northern IVCs. Like these clouds, the IVC may be part of a Galactic fountain flow, but it is associated with the Perseus arm rather than with the Local arm.

4. We determine the distance to HVC Complex GCP as 9.8 to 15.1 kpc, based on the detection of Ca II K absorption ($51 \pm 3 \text{ m}\text{\AA}$) in the spectrum of the RR Lyrae star V1084 Aql ($D = 14.5 \pm 1.3 \text{ kpc}$) and the significant nondetection (3σ limit $11 \text{ m}\text{\AA}$, expected $59 \text{ m}\text{\AA}$) toward the RR Lyrae V1172 Aql ($D = 10.5 \pm 1.4 \text{ kpc}$).

5. The distance bracket for Complex GCP implies that the cloud is at a galactocentric radius of 6.2 to 9.7 kpc, lies 2.5 to 3.9 kpc below the plane, and has a neutral gas mass of $\sim 10^6 M_{\odot}$. $\text{H}\alpha$ emission is detected from two positions in the cloud and implies an ionized component that is 40% to 120% as large as the neutral component. At its location the cloud is moving with a velocity of $35\text{--}75 \text{ km s}^{-1}$ relative to gas taking part in differential galactic rotation, and it is moving away from the Galactic plane at about $10\text{--}20 \text{ km s}^{-1}$.

6. For the IVC cloud g1 we derive a distance bracket of 1.8 to 3.8 kpc, detecting the associated absorption in the spectra of

three stars, 2MASS J213327.04+133026.7 ($D = 8.2 \pm 0.9$ kpc), FW Peg ($D = 3.6 \pm 0.8$ kpc), and BS 17578–0016 ($D = 3.6 \pm 0.5$), but not in four nearer stars. This places the IVC 0.8 to 1.7 kpc below the Galactic disk and implies that it has a peculiar velocity of about 55 km s^{-1} . It is a small cloud, with a mass of no more than $10^4 M_{\odot}$ and an area of $(60\text{--}130)^2 \text{ pc}^2$. It is moving away from the Galactic plane with a velocity of about 20 km s^{-1} . The relatively bright $\text{H}\alpha$ emission coming from this cloud (Smoker et al. 2002) implies that there is a factor of 5 to 10 more ionized gas than neutral gas. The most likely interpretation for this object appears to be that it is condensing in an upward fountain flow.

7. The $\text{Ca II}/\text{H I}$ ratios that we derive are all in line with the relation between $R(\text{Ca II})$ and $N(\text{H I})$ that was found by Wakker & Mathis (2000), except that each cloud appears to have a different scaling factor. In particular, the Cohen Stream detections are a factor of 0.44 ± 0.5 below the average relation, for the Perseus arm gas the scaling factor is 0.62 ± 0.11 , the Complex GCP detection is on the average, while the cloud g1 detections fall a factor of 2.2 ± 1.0 above the average.

8. We have set new distance brackets to four HVCs/IVCs, which trace at least two of the different origins of HVCs: inflow of new material (the Cohen Stream) and the Galactic fountain (cloud g1 and the Perseus arm IVC). The origin of the fourth cloud (Complex GCP) is less clear, but with our distance estimate detailed modeling is now possible. In the future we hope to determine more distances to HVCs/IVCs, and we should eventually be able to measure the inflow rate of accreting material, as well as the rate of circulation of gas between disk and halo.

B. P. W., D. G. Y., R. W., and T. C. B. acknowledge support from grant AST 06-07154 awarded by the US National Science Foundation. T. C. B. also acknowledges NSF grant PHY-02-16783, Physics Frontier Center/Joint Institute for Nuclear Astrophysics (JINA).

We thank Ben McCall for help in observing logistics support, the staff of Apache Point Observatory for excellent observing support, and Chris Thom for discussions.

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the US Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max-Planck Society, and the Higher Education Funding Council for England. The SDSS Web site is <http://www.sdss.org/>.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.

REFERENCES

- Adelman-McCarthy, J., et al. 2007, *ApJS*, 172, 634
 Albert, C. E., Blades, J. C., Morton, D. C., Lockman, F. J., Proulx, M., & Ferrarese, L. 1993, *ApJS*, 88, 81
 Beers, T. C., Wilhelm, R., Doinidis, S. P., & Mattson, C. J. 1996, *ApJS*, 103, 433
 Bland-Hawthorn, J., & Maloney, P. R. 1999, *ApJ*, 510, L33
 Bland-Hawthorn, J., Veilleux, S., Cecil, G. N., Putman, M. E., Gibson, B. K., & Maloney, P. R. 1998, *MNRAS*, 299, 611
 Bond, H. E. 1978, *PASP*, 90, 526
 Bowen, D. V., Roth, K. C., Meyer, D. M., & Blades, J. C. 2000, *ApJ*, 536, 225
 Brown, W. R., Geller, M. J., Kenyon, S. J., Beers, T. C., Kurtz, M. J., & Roll, J. B. 2004, *AJ*, 127, 1555
 Burton, W. B., & te Lintel Hekkert, P. 1986, *A&AS*, 65, 427
 Chiappini, C., Matteucci, F., & Romano, D. 2001, *ApJ*, 554, 1044
 Cohen, J. G. 1979, *ApJ*, 231, 751
 Cohen, R. J. 1981, *MNRAS*, 196, 835
 d'Odorico, S., di Serego Alighieri, S., Pettini, M., Magain, P., Nissen, P. E., & Panagia, N. 1989, *A&A*, 215, 21
 d'Odorico, S., Pettini, M., & Ponz, D. 1985, *ApJ*, 299, 852
 Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
 Gibson, B. K., Giroux, M. L., Penton, S. V., Putman, M. E., Stocke, J. T., & Shull, J. M. 2000, *AJ*, 120, 1830
 Girardi, L., Grebel, E. K., Odenkirchen, M., & Chiosi, C. 2004, *A&A*, 422, 205
 Green, R. F., Schmidt, M., & Liebert, J. 1986, *ApJS*, 61, 305
 Gunn, J. E., et al. 1998, *AJ*, 116, 3040
 ———. 2006, *AJ*, 131, 2332
 Haffner, L. M., Reynolds, R. J., & Tufte, S. L. 1999, *ApJ*, 523, 223
 Haffner, L. M., Reynolds, R. J., Tufte, S. L., Madsen, G. H., Jaehnig, K. P., & Percival, J. W. 2003, *ApJS*, 149, 405
 Hartmann, D., & Burton, W. B. 1997, *Atlas of Galactic Neutral Hydrogen* (Cambridge: Cambridge Univ. Press)
 Ho, L. C., & Filippenko, A. V. 1995, *ApJ*, 444, 165
 ———. 1996, *ApJ*, 463, 818
 Hogg, D. W., Finkbeiner, D. P., Schlegel, D. J., & Gunn, J. E. 2001, *AJ*, 122, 2129
 Houck, J. C., & Bregman, J. N. 1990, *ApJ*, 352, 506
 Hulsbosch, A. N. M., & Wakker, B. P. 1988, *A&AS*, 75, 191
 Ivezić, Ž., Vivas, A. K., Lupton, R. H., & Zinn, R. 2005, *AJ*, 129, 1096
 Ivezić, Ž., et al. 2000, *AJ*, 120, 963
 ———. 2004, *Astron. Nachr.*, 325, 583
 Kalberla, P. W. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, *A&A*, 440, 775
 Kemp, S. N., Bates, B., Dufton, P. L., Keenan, F. P., & Montgomery, A. S. 1994, *MNRAS*, 270, 597
 Kennedy, D. C., Bates, B., Keenan, F. P., Kemp, S. N., Ryans, R. S. I., Davies, R. D., & Sembach, K. R. 1998, *MNRAS*, 297, 849
 Kukarkin, B. V., et al. 1969, *General Catalogue of Variable Stars*, Vol. 1 (3rd ed.; Moscow: Sternberg Institute)
 ———. 1970, *General Catalogue of Variable Stars*, Vol. 2 (3rd ed.; Moscow: Sternberg Institute)
 Kuntz, K. D., & Danly, L. 1996, *ApJ*, 457, 703
 Kutyrev, A. S., & Reynolds, R. J. 1989, *ApJ*, 344, L9
 Layden, A. C., Hanson, R. D., Hawley, S. L., Klemola, A. R., & Hanley, C. J. 1996, *AJ*, 112, 2110
 Lehner, N., Rolleston, W. R. J., Ryans, R. S. I., Keenan, F. P., Bates, B., Pollacco, D. L., & Sembach, K. R. 1999, *A&AS*, 134, 257
 Little, J. E., Dufton, P. L., Keenan, F. P., & Conlon, E. S. 1994, *ApJ*, 427, 267
 Lu, L., Savage, B. D., Sembach, K. R., Wakker, B. P., Sargent, W. W. L., & Oosterloo, T. A. 1998, *AJ*, 115, 162
 Meyer, D. M., & Roth, K. C. 1991, *ApJ*, 383, L41
 Moehler, S., Heber, U., & de Boer, K. S. 1990, *A&A*, 239, 265
 Morras, R., Bajaja, E., Arnal, E. M., & Pöppel, W. G. L. 2000, *A&AS*, 142, 25
 Pier, J. R., Munn, J. A., Hindsley, R. B., Hennessy, G. S., Kent, S. M., Lupton, R. H., & Ivezić, Ž. 2003, *AJ*, 125, 1559
 Phillips, J. P. 2004, *MNRAS*, 353, 589
 Prata, S., & Wallerstein, G. 1967, *PASP*, 79, 202
 Preston, G. W., Schectman, S. A., & Beers, T. C. 1991, *ApJS*, 76, 1001
 Putman, M. E., Thom, C., Gibson, B. K., & Staveley-Smith, L. 2004, *ApJ*, 603, L77
 Reynolds, R. J., Tufte, S. L., Kung, D. T., McCullough, P. R., & Heiles, C. 1995, *ApJ*, 448, 715
 Richter, P. 2006, *Rev. Mod. Astron.*, 19, 31
 Richter, P., Savage, B. D., Wakker, B. P., Sembach, K. R., & Kalberla, P. M. W. 2001a, *ApJ*, 549, 281
 Richter, P., Sembach, K. R., Wakker, B. P., Savage, B. D., Tripp, T. M., Murphy, E. M., Kalberla, P. M. W., & Jenkins, E. B. 2001b, *ApJ*, 559, 318

- Robertson, J. G., Schwarz, U. J., van Woerden, H., Murray, J. D., Morton, D. C., & Hulsbosch, A. N. M. 1991, *MNRAS*, 248, 508
- Ryans, R. S. I., Sembach, K. R., & Keenan, F. P. 1996, *A&A*, 314, 609
- Savage, B. D., & Sembach, K. R. 1991, *ApJ*, 379, 245
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schwarz, U. J., & Wakker, B. P. 2004, in *High Velocity Clouds*, ed. H. van Woerden et al. (Dordrecht: Kluwer), 145
- Schwarz, U. J., Wakker, B. P., & van Woerden, H. 1995, *A&A*, 302, 364
- Sembach, K. R., Savage, B. D., & Massa, D. 1991, *ApJ*, 372, 81
- Sembach, K. R., et al. 2004, *ApJS*, 150, 387
- Sirko, E., et al. 2004, *AJ*, 127, 899
- Smith, G. P. 1963, *Bull. Astron. Inst. Netherlands*, 17, 203
- Smith, J. A., et al. 2002, *AJ*, 123, 2121
- Smoker, J. V., Haffner, L. M., Keenan, F. P., Davies, R. D., & Pollacco, D. 2002, *MNRAS*, 337, 385
- Smoker, J. V., Lehner, N., Keenan, F. P., Totten, E. J., Murphy, E., Sembach, K. R., Davies, R. D., & Bates, B. 2001, *MNRAS*, 322, 13
- Songaila, A., & York, D. G. 1980, *ApJ*, 242, 976
- Stoughton, C., et al. 2002, *AJ*, 123, 485
- Tamanaha, C. M. 1995, *ApJ*, 450, 638
- . 1996, *ApJS*, 104, 81
- Thom, C., Putman, M. E., Gibson, B. K., Christlieb, N., Flynn, C., Beers, T. C., Wilhelm, R., & Lee, Y. S. 2006, *ApJ*, 638, L97
- Tripp, T. M., et al. 2003, *AJ*, 125, 3122
- Tucker, D., et al. 2006, *Astron. Nachr.*, 327, 821
- van Woerden, H., Schwarz, U. J., Peletier, R. F., Wakker, B. P., & Kalberla, P. M. W. 1999, *Nature*, 400, 138
- van Woerden, H., & Wakker, B. P. 2004, in *High Velocity Clouds*, ed. H. van Woerden et al. (Dordrecht: Kluwer), 195
- Vladilo, G., Centurión, M., de Boer, K. S., King, D. L., Lipman, K., Stegert, J. S. W., Unger, S. W., & Walton, N. A. 1994, *A&A*, 291, 425
- Wakker, B. P. 1991, *A&A*, 250, 499
- . 2001, *ApJS*, 136, 463
- . 2004, in *High Velocity Clouds*, ed. H. van Woerden et al. (Dordrecht: Kluwer), 25
- Wakker, B. P., Kalberla, P. M. W., van Woerden, H., de Boer, K. S., & Putman, M. E. 2001, *ApJS*, 136, 537
- Wakker, B. P., & Mathis, J. S. 2000, *ApJ*, 544, L107
- Wakker, B. P., & van Woerden, H. 1991, *A&A*, 250, 509
- . 1997, *ARA&A*, 35, 217
- Wakker, B. P., van Woerden, H., Schwarz, U. J., Peletier, R. F., & Douglas, N. G. 1996, *A&A*, 306, L25
- Wakker, B. P., et al. 1999, *Nature*, 402, 388
- . 2003, *ApJS*, 146, 1
- Weiner, B. J., Vogel, S. N., & Williams, T. B. 2001, in *ASP Conf. Ser. 240, Gas and Galaxy Evolution*, ed. J. E. Hibbard, M. P. Rupen, & J. H. van Gorkom (San Francisco: ASP), 515
- West, K. A., Pettini, M., Penton, M., Blades, J. C., & Morton, D. C. 1985, *MNRAS*, 215, 481
- Wilhelm, R., Beers, T. C., & Gray, R. O. 1999, *AJ*, 117, 2308
- York, D. G., et al. 2000, *AJ*, 120, 1579